

GLOBAL WELL-POSEDNESS FOR THE COMPRESSIBLE NAVIER-STOKES EQUATIONS WITH THE HIGHLY OSCILLATING INITIAL VELOCITY

QIONGLEI CHEN, CHANGXING MIAO, AND ZHIFEI ZHANG

ABSTRACT. Cannone [3] proved the global well-posedness of the incompressible Navier-Stokes equations for a class of highly oscillating data. In this paper, we prove the global well-posedness for the compressible Navier-Stokes equations in the critical functional framework with the initial data close to a stable equilibrium. Especially, this result allows us to construct global solutions for the highly oscillating initial velocity. The proof relies on a new estimate for the hyperbolic/parabolic system with convection terms.

1. INTRODUCTION

We consider the compressible Navier-Stokes equations in $\mathbb{R}^+ \times \mathbb{R}^n (n = 2, 3)$

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \nabla P(\rho) = 0, \\ (\rho, u)|_{t=0} = (\rho_0, u_0). \end{cases} \quad (1.1)$$

Here $\rho(t, x)$ and $u(t, x)$ denote the density and velocity of the fluid respectively. The pressure P is a suitably smooth function of ρ , and the Lamé coefficients μ and λ satisfy

$$\mu > 0 \quad \text{and} \quad \lambda + 2\mu > 0, \quad (1.2)$$

which ensures that the operator $\mu \Delta + (\lambda + \mu) \nabla \operatorname{div}$ is elliptic.

The local existence and uniqueness of smooth solutions for the system (1.1) were proved by Nash [24] for smooth initial data without vacuum. Later on, Matsumura and Nishida [23] proved the global well-posedness for smooth data close to equilibrium. Recently, Lions [22] proved the global existence of weak solutions for large initial data. However, the question of uniqueness of weak solutions remains open, even in the two dimensional case.

A natural way of dealing with the problem of uniqueness is to find a functional setting as large as possible in which existence and uniqueness hold. For the incompressible Navier-Stokes equations

$$\begin{cases} \partial_t u - \nu \Delta u + u \cdot \nabla u + \nabla p = 0, \\ \operatorname{div} u = 0, \\ u(x, 0) = u_0, \end{cases} \quad (1.3)$$

such an approach was initiated by Fujita and Kato [17]. They proved the global existence and uniqueness of strong solution for small initial data in the homogeneous

Date: 2, August, 2009.

2000 Mathematics Subject Classification. 35Q30, 35B35.

Key words and phrases. Compressible Navier-Stokes equations, Global well-posedness, Besov space, highly oscillating.

Sobolev space $\dot{H}^{\frac{n}{2}-1}$. Let's point out that the space $\dot{H}^{\frac{n}{2}-1}$ is a critical space. We give the precise meaning of critical spaces. It is well-known that if (u, p) is a solution of (1.3), then

$$(u_\lambda(t, x), p_\lambda(t, x)) \stackrel{\text{def}}{=} (\lambda u(\lambda^2 t, \lambda x), \lambda^2 p(\lambda^2 t, \lambda x)) \quad (1.4)$$

is also a solution of (1.3). A functional space is called critical if the corresponding norm is invariant under the scaling of (1.4). For the compressible Navier-Stokes equations, one can check that if (ρ, u) is a solution of (1.1), then

$$(\rho_\lambda(t, x), u_\lambda(t, x)) \stackrel{\text{def}}{=} (\rho(\lambda^2 t, \lambda x), \lambda u(\lambda^2 t, \lambda x)),$$

is also a solution of (1.1) provided the pressure law has been changed into $\lambda^2 P$. This motivates the following definition.

Definition 1.1. A functional space is called critical if the associated norm is invariant under the transformation $(\rho, u) \longrightarrow (\rho_\lambda, u_\lambda)$ (up to a constant independent of λ).

Consequently, a natural candidate is the homogenous Sobolev space $\dot{H}^{\frac{n}{2}} \times (\dot{H}^{\frac{n}{2}-1})^n$. However, $\dot{H}^{\frac{n}{2}}$ is not included in L^∞ such that we cannot expect to obtain a L^∞ control of the density when $\rho_0 - \bar{\rho} \in \dot{H}^{\frac{n}{2}}$. Instead, one can choose the initial data (ρ_0, u_0) such that for some $\bar{\rho}$,

$$(\rho_0 - \bar{\rho}, u_0) \in \dot{B}_{2,1}^{\frac{n}{2}} \times (\dot{B}_{2,1}^{\frac{n}{2}-1})^n,$$

since $\dot{B}_{2,1}^{\frac{n}{2}}$ is continuously embedded in L^∞ . In a seminal paper, Danchin[11] proved the global well-posedness of (1.1) in the critical Besov space for small initial data u_0 and ρ_0 close to $\bar{\rho}$. More precisely,

Theorem 1.2. Let $\bar{\rho} > 0$ be such that $P'(\bar{\rho}) > 0$. There exist two positive constants c and M such that for all (ρ_0, u_0) with $\rho_0 - \bar{\rho} \in \dot{B}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}$, $u_0 \in \dot{B}_{2,1}^{\frac{n}{2}-1}$ and

$$\|\rho_0 - \bar{\rho}\|_{\dot{B}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}} + \|u_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq c$$

then the system (1.1) has a global unique solution $(\rho - \bar{\rho}, u)$ such that

$$\|(\rho_0 - \bar{\rho}, u)\|_{F^{\frac{n}{2}}} \leq M(\|\rho_0 - \bar{\rho}\|_{\dot{B}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}} + \|u_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}),$$

where

$$F^s \stackrel{\text{def}}{=} L^1(\mathbb{R}^+; \dot{B}^{s+1, s}) \cap C(\mathbb{R}^+; \dot{B}^{s-1, s}) \times \left(L^1(\mathbb{R}^+; \dot{B}_{2,1}^{s+1}) \cap C(\mathbb{R}^+; \dot{B}_{2,1}^{s-1}) \right)^n.$$

Here and what follows, we refer to Section 2.2 for the definition of Besov spaces.

On the other hand, Fujita-Kato result was generalized to the critical Lebesgue space L^n by Weissler [26](see also [21]), and to a class of function spaces such as $\dot{B}_{n,\infty}^0$ by Cannone, Meyer and Planchon [4] which allows to construct self-similar solutions. Recently, Cannone [3] generalized it to Besov spaces of negative index of regularity. More precisely, he showed that if the initial data satisfies

$$\|u_0\|_{\dot{B}_{p,\infty}^{-1+\frac{n}{p}}} \leq c, \quad p > n$$

for some small constant c , then the Navier-Stokes equations (1.3) is globally well-posed. Let us emphasize that this result allows us to construct global solutions for

highly oscillating initial data which may have a large norm in $\dot{H}^{\frac{n}{2}-1}$ or L^n . A typical example in three dimensions is

$$u_0(x) = \sin\left(\frac{x_3}{\varepsilon}\right)(-\partial_2\phi(x), \partial_1\phi(x), 0)$$

where $\phi \in \mathcal{S}(\mathbb{R}^3)$ and $\varepsilon > 0$ is small enough. We refer to [8, 10, 9, 25] for some relevant results. An important question is then to prove a theorem of this type for the compressible Navier-Stokes equations.

To solve this problem, the key point is to generalize the well-posed result in Theorem 1.2 to the critical functional spaces with negative index of regularity (the corresponding norm of the highly oscillating function is small). A typical space is $\dot{B}_{p,1}^{\frac{n}{p}} \times (\dot{B}_{p,1}^{\frac{n}{p}-1})^n$ for some $p > n$. Let's recall the proof of Theorem 1.2, where the energy method is applied to obtain the smoothing effect and damping effect of the system. This method works well for the well-posed problem in the L^2 framework. However, it doesn't work for the well-posed problem in the L^p framework for $p \neq 2$, especially in the case when the coupling effect of the system is important for the problem. In this paper, we will develop a new method based on Green's matrix of the linearized system to study the well-posed problem in general Besov spaces.

In order to present our method, let us look at the linearized system of (1.1) with the convection terms:

$$\begin{cases} \partial_t a + \Lambda d = -v \cdot \nabla a + F, \\ \partial_t d - \bar{v} \Delta d - \Lambda a = -v \cdot \nabla d + G, \\ (a, d)|_{t=0} = (a_0, d_0), \end{cases} \quad (1.5)$$

where $\Lambda = (-\Delta)^{\frac{1}{2}}$. Let $\mathcal{G}(x, t)$ be the Green matrix of the linear system

$$\begin{cases} \partial_t a + \Lambda d = 0, \\ \partial_t d - \bar{v} \Delta d - \Lambda a = 0. \end{cases}$$

Then the solution (a, d) of (1.5) can be expressed as

$$\begin{pmatrix} a \\ d \end{pmatrix} = \mathcal{G}(x, t) * \begin{pmatrix} a_0 \\ d_0 \end{pmatrix} + \int_0^t \mathcal{G}(x, t - \tau) * \begin{pmatrix} F - v \cdot \nabla a \\ G - v \cdot \nabla d \end{pmatrix} d\tau. \quad (1.6)$$

To obtain the estimates of (a, d) in the Besov spaces, it is natural to study the action of Green's matrix $\mathcal{G}(x, t)$ on distributions whose Fourier transform is supported in a ring. As we know, the semi-group of the heat equation shows for any couple (t, λ) of positive real numbers,

$$\text{supp } \hat{f} \subset \lambda \mathcal{C} \implies \|e^{t\Delta} f\|_{L^p} \leq C e^{-ct\lambda^2} \|f\|_{L^p}, \quad 1 \leq p \leq \infty,$$

where $\mathcal{C}$ is a ring, which plays an important role in a new proof given by Chemin[7] on the Cannone, Meyer and Planchon result. To obtain similar properties for $\mathcal{G}(x, t)$, we need to investigate the precise behaviour of $\widehat{\mathcal{G}}(\xi, t)$. Roughly speaking, for $|\xi| \lesssim 1$, $\widehat{\mathcal{G}}(\xi, t)$ behaves like the heat kernel:

$$|D_\xi^\alpha \widehat{\mathcal{G}}(\xi, t)| \leq C e^{-\vartheta|\xi|^2 t} (1 + |\xi|)^{|\alpha|} (1 + t)^{|\alpha|},$$

which allows us to obtain

$$\|\mathcal{G} * f\|_{L^2} \leq C e^{-ct\lambda^2} \|f\|_{L^2}, \quad (1.7)$$

if $\text{supp } \hat{f} \subset \lambda \mathcal{C}$ for $\lambda \lesssim 1$. For $|\xi| \gg 1$, $\widehat{\mathcal{G}}(\xi, t)$ has the following precise expansion:

$$\begin{aligned} \widehat{\mathcal{G}}(\xi, t) = & e^{-\bar{\nu}^{-1}t} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + e^{-\bar{\nu}|\xi|^2 t} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ & + \widehat{\mathcal{G}}^1(\xi, t) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \widehat{\mathcal{G}}^2(\xi, t) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \end{aligned}$$

where $\widehat{\mathcal{G}}^1$ and $\widehat{\mathcal{G}}^2$ satisfy the estimates

$$\begin{aligned} |\partial_\xi^\alpha \widehat{\mathcal{G}}^1| & \leq C|\xi|^{-|\alpha|-1} (e^{-\frac{\bar{\nu}^{-1}t}{2}} + e^{-\vartheta|\xi|^2 t}), \\ |\partial_\xi^\alpha \widehat{\mathcal{G}}^2| & \leq C|\xi|^{-|\alpha|-2} (e^{-\frac{\bar{\nu}^{-1}t}{2}} + e^{-\vartheta|\xi|^2 t}), \end{aligned}$$

which allow us to obtain, for any $1 \leq p \leq \infty$,

$$\|\mathcal{G}^1 * f\|_{L^p} \leq C\lambda^{-1}e^{-ct}\|f\|_{L^p}, \quad \|\mathcal{G}^2 * f\|_{L^p} \leq C\lambda^{-2}e^{-ct}\|f\|_{L^p}, \quad (1.8)$$

if $\text{supp } \hat{f} \subset \lambda \mathcal{C}$ for $\lambda \gg 1$. Since $\mathcal{G}$ has different behaviour for low frequency and high frequency, it is natural to work with the hybrid-Besov spaces (see Definition 2.6). Indeed, from (1.6), (1.7) and (1.8), we can obtain the following estimate in the hybrid-Besov spaces: for any $1 \leq r \leq \infty$,

$$\begin{aligned} & \|a\|_{\widetilde{L}_T^r(\dot{\mathcal{B}}_{2,p}^{s_p-1+\frac{2}{r},s})} + \|d\|_{\widetilde{L}_T^r(\dot{\mathcal{B}}_{2,p}^{s_p-1+\frac{2}{r},s-1+\frac{2}{r}})} \\ & \leq C \left[\|a_0\|_{\dot{\mathcal{B}}_{2,p}^{s_p-1,s}} + \|d_0\|_{\dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}} + \|F\|_{L_T^1(\dot{\mathcal{B}}_{2,p}^{s_p-1,s})} + \|G\|_{L_T^1(\dot{\mathcal{B}}_{2,p}^{s_p-1,s-1})} \right. \\ & \quad \left. + \|v \cdot \nabla a\|_{L_T^1(\dot{\mathcal{B}}_{2,p}^{s_p-1,s})} + \|v \cdot \nabla d\|_{L_T^1(\dot{\mathcal{B}}_{2,p}^{s_p-1,s-1})} \right], \quad (1.9) \end{aligned}$$

where $s_p \stackrel{\text{def}}{=} s - \frac{n}{p} + \frac{n}{2}$. Let us point out that if we take $s = \frac{n}{p}$ for $p > n$, the regularity index $s - 1$ for the high frequency part of d_0 is negative, which is enough to ensure that the norm of d_0 in the hybrid-Besov space $\dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}$ is small if d_0 is highly oscillating, see Proposition 2.9.

Unfortunately, the inequality (1.9) cannot be applied directly. In fact, if the convection term $v \cdot \nabla a$ is viewed as a perturbation term, one derivative loss about the function a will appear no matter how smooth is v . To overcome this difficulty, we will work in the Lagrangian coordinate such that the convection terms in (1.5) will disappear modulus some commutators. Here the idea is partially motivated by the recent work [18, 19] for the incompressible flows.

We denote

$$\begin{aligned} \mathcal{E}^s \stackrel{\text{def}}{=} & \left\{ (a, d) \in \left(L^1(\mathbb{R}^+; \dot{\mathcal{B}}_{2,p}^{s_p+1,s}) \cap C(\mathbb{R}^+; \dot{\mathcal{B}}_{2,p}^{s_p-1,s}) \right) \right. \\ & \left. \times \left(L^1(\mathbb{R}^+; \dot{\mathcal{B}}_{2,p}^{s_p+1,s+1}) \cap C(\mathbb{R}^+; \dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}) \right)^n \right\} \end{aligned}$$

with $s_p \stackrel{\text{def}}{=} s - \frac{n}{p} + \frac{n}{2}$. Our main result is stated as follows.

Theorem 1.3. Let $\bar{\rho}$ be a positive constant such that $P'(\bar{\rho}) > 0$. There exist two positive constants c and M such that for all (ρ_0, u_0) with $\rho_0 - \bar{\rho} \in \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}$, $u_0 \in \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}$ and

$$\|\rho_0 - \bar{\rho}\|_{\dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} + \|u_0\|_{\dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \leq \eta, \quad (1.10)$$

the following results hold

(1) **Existence.** if $2 \leq p < 2n$, and $p \leq \min(4, \frac{2n}{n-2})$, the system (1.1) has a global solution $(\rho - \bar{\rho}, u) \in \mathcal{E}^{\frac{n}{p}}$ with

$$\|(\rho - \bar{\rho}, u)\|_{\mathcal{E}^{\frac{n}{p}}} \leq M(\|\rho_0 - \bar{\rho}\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} + \|u_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}).$$

(2) **Uniqueness.** if $2 \leq p \leq n$, then the uniqueness holds in $\mathcal{E}^{\frac{n}{p}}$.

Remark 1.4. Compared with Theorem 1.2, an important improvement of Theorem 1.3 is that it allows the regularity index for the high frequency part of u_0 to be negative. Especially, this allows us to obtain the global well-posedness of (1.1) for the highly oscillating initial velocity u_0 . For example,

$$u_0(x) = \sin\left(\frac{x_1}{\varepsilon}\right)\phi(x), \quad \phi(x) \in \mathcal{S}(\mathbb{R}^n),$$

which satisfies

$$\|u_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \ll 1 \quad \text{for } p > n$$

if ε is small enough, see Proposition 2.9.

Remark 1.5. In fact, Theorem 1.3 also holds for general dimension. But we need to restrict the dimension $n = 2, 3$ in order to obtain global solutions for the highly oscillating initial velocity.

Remark 1.6. We believe that our method can be adapted to the other hyperbolic-parabolic models. This is the object of our future work.

The structure of this paper is as follows.

In Section 2, we recall some basic facts about Littlewood-Paley theory and the functional spaces. In Section 3, we recall some results concerning the classical parabolic regularizing effect. Section 4 is devoted to the study of Green's matrix of the linearized system. In Section 5, we establish the key estimates for the linearized system with convection terms. Section 6 is devoted to the proof of Theorem 1.3. Finally, an appendix is devoted to some estimates in the hybrid-Besov spaces.

2. LITTLEWOOD-PALEY THEORY AND THE FUNCTION SPACES

2.1. Littlewood-Paley theory. First of all, we introduce the Littlewood-Paley decomposition. Choose two radial functions $\varphi, \chi \in \mathcal{S}(\mathbb{R}^n)$ supported in $\mathcal{C} = \{\xi \in \mathbb{R}^n, \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$, $\mathcal{B} = \{\xi \in \mathbb{R}^n, |\xi| \leq \frac{4}{3}\}$ respectively such that

$$\sum_{j \in \mathbb{Z}} \varphi(2^{-j}\xi) = 1 \quad \text{for all } \xi \neq 0.$$

The frequency localization operators Δ_j and S_j are defined by

$$\Delta_j f = \varphi(2^{-j}D)f, \quad S_j f = \sum_{k \leq j-1} \Delta_k f = \chi(2^{-j}D)f, \quad \text{for } j \in \mathbb{Z}.$$

With our choice of φ , it is easy to verify that

$$\begin{aligned} \Delta_j \Delta_k f &= 0 \quad \text{if } |j - k| \geq 2 \quad \text{and} \\ \Delta_j (S_{k-1} f \Delta_k f) &= 0 \quad \text{if } |j - k| \geq 5. \end{aligned} \tag{2.1}$$

In the sequel, we will constantly use the Bony's decomposition from [2]:

$$fg = T_fg + T_gf + R(f, g), \quad (2.2)$$

with

$$T_fg = \sum_{j \in \mathbb{Z}} S_{j-1}f \Delta_j g, \quad R(f, g) = \sum_{j \in \mathbb{Z}} \Delta_j f \tilde{\Delta}_j g, \quad \tilde{\Delta}_j g = \sum_{|j'-j| \leq 1} \Delta_{j'} g.$$

Next, let us introduce some useful lemmas which will be repeatedly used throughout this paper.

Lemma 2.1. [6] Let $1 \leq p \leq q \leq +\infty$. Then for any $\gamma \in (\mathbb{N} \cup \{0\})^n$, there exists a constant C independent of f, j such that

$$\begin{aligned} \text{supp} \hat{f} \subseteq \{|\xi| \leq A_0 2^j\} &\Rightarrow \|\partial^\gamma f\|_{L^q} \leq C 2^{j|\gamma| + jn(\frac{1}{p} - \frac{1}{q})} \|f\|_{L^p}, \\ \text{supp} \hat{f} \subseteq \{A_1 2^j \leq |\xi| \leq A_2 2^j\} &\Rightarrow \|f\|_{L^p} \leq C 2^{-j|\gamma|} \sup_{|\beta|=|\gamma|} \|\partial^\beta f\|_{L^p}. \end{aligned}$$

Let a vector field $v(t, x) \in L^1_{\text{loc}}(\mathbb{R}; \text{Lip}(\mathbb{R}^n))$. We define ψ and ψ_j as follows

$$\begin{aligned} \frac{d}{dt} \psi(t, x) &= v(t, \psi(t, x)), \quad \psi(0, x) = x, \\ \frac{d}{dt} \psi_j(t, x) &= S_{j-1} v(t, \psi_j(t, x)), \quad \psi_j(0, x) = x, \quad j \in \mathbb{Z}. \end{aligned}$$

In what follows, we denote $V(t) \stackrel{\text{def}}{=} \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau$. Now we are position to state the following lemmas.

Lemma 2.2. Let $1 \leq p < \infty$. Then there hold

$$\|f \circ \psi(t, x)\|_{L^p} \leq e^{V(t)} \|f\|_{L^p}, \quad \|f \circ \psi(t, x)\|_{L^\infty} \leq \|f\|_{L^\infty}.$$

Proof. Lemma 2.2 can be deduced from the fact

$$\partial_t \det(\nabla \psi) = \text{div } v \det(\nabla \psi),$$

and a change of variables. □

Lemma 2.3. Let $1 \leq p \leq \infty$. Then for any $j, k \in \mathbb{Z}$, there hold

$$\begin{aligned} \|\Delta_j(\Delta_k f \circ \psi_k)\|_{L^p} &\leq C 2^{-(j-k)} e^{CV(t)} \|\Delta_k f\|_{L^p}, \\ \|S_j(\Delta_k f \circ \psi_k)\|_{L^p} &\leq C e^{CV(t)} (V(t) + 2^{j-k}) \|\Delta_k f\|_{L^p}. \end{aligned}$$

Here C is a constant independent of j, k .

Proof. Let $\{\alpha_\ell\}_{\ell=1}^n$ be a smooth unity decomposition of $\mathbb{S}^{n-1}$ such that $\xi'_\ell \neq 0$ on $\text{supp } \alpha_\ell(\xi')$, here $\xi' = (\xi'_1, \dots, \xi'_n) \in \mathbb{S}^{n-1}$. Then we write

$$\varphi(\xi) = \sum_{\ell=1}^n i \xi_\ell \hat{\theta}_\ell(\xi), \quad \hat{\theta}_\ell(\xi) \stackrel{\text{def}}{=} (i \xi_\ell)^{-1} \alpha_\ell(\xi/|\xi|) \varphi(\xi),$$

consequently,

$$\Delta_j f = 2^{-j} \sum_{\ell=1}^n \partial_\ell (2^{jn} \theta_\ell(2^j \cdot) * f) \stackrel{\text{def}}{=} 2^{-j} \sum_{\ell=1}^n \partial_\ell (\bar{\Delta}_{j\ell} f), \quad j \in \mathbb{Z}, \quad (2.3)$$

from which, we get by integration by parts that

$$\Delta_j(\Delta_k f \circ \psi_k) = \sum_{\ell=1}^n 2^{nj-j} \int_{\mathbb{R}^n} \theta_\ell(2^j(x-y)) \partial_\ell((\Delta_k f)(\psi_k(y))) dy.$$

Thus by Young's inequality, Lemma 2.2 and Lemma 2.1, we have

$$\begin{aligned} \|\Delta_j(\Delta_k f \circ \psi_k)\|_{L^p} &\leq C 2^{-j} \|(\nabla \Delta_k f)(\psi_k)\|_{L^p} \|\nabla \psi_k\|_{L^\infty} \\ &\leq C 2^{-j} e^{V(t)} \|\nabla \Delta_k f\|_{L^p} \|\nabla \psi_k\|_{L^\infty} \\ &\leq C 2^{k-j} e^{V(t)} \|\Delta_k f\|_{L^p} \|\nabla \psi_k\|_{L^\infty}. \end{aligned} \quad (2.4)$$

On the other hand, thanks to (2.3), we have

$$S_j(\Delta_k f \circ \psi_k) = 2^{-k} \sum_{\ell=1}^n S_j(\partial_\ell(\bar{\Delta}_{k\ell} f) \circ \psi_k).$$

Set $g(x) \stackrel{\text{def}}{=} (\mathcal{F}^{-1} \chi)(x)$. Making a change of variables: $y \mapsto \psi_k^{-1}(y)$, and then integrating by parts yields

$$\begin{aligned} S_j(\Delta_k f \circ \psi_k) &= 2^{nj-k} \sum_{\ell=1}^n \int_{\mathbb{R}^n} g(2^j(x-y)) (\partial_\ell(\bar{\Delta}_{k\ell} f)(\psi_k(y))) dy \\ &= 2^{nj-k} \sum_{\ell=1}^n \int_{\mathbb{R}^n} g(2^j(x - \psi_k^{-1}(y))) \partial_\ell(\bar{\Delta}_{k\ell} f)(y) \det(\nabla \psi_k^{-1}) dy \\ &= 2^{nj-k} \sum_{\ell=1}^n \int_{\mathbb{R}^n} \left\{ 2^j \partial_t \psi_k^{-1}(\partial_{y_\ell} g)(2^j(x - \psi_k^{-1}(y))) \det(\nabla \psi_k^{-1}) \right. \\ &\quad \left. + g(2^j(x - \psi_k^{-1}(y))) \partial_\ell \det(\nabla \psi_k^{-1}) \right\} \bar{\Delta}_{k\ell} f(y) dy. \end{aligned}$$

Thus, we get by Young's inequality that

$$\begin{aligned} \|S_j(\Delta_k f \circ \psi_k)\|_{L^p} &\leq C 2^{j-k} \|\nabla \psi_k^{-1}\|_{L^\infty}^{n+1} \|\Delta_k f\|_{L^p} \\ &\quad + C 2^{-k} \|\nabla^2 \psi_k^{-1}\|_{L^\infty} \|\nabla \psi_k^{-1}\|_{L^\infty}^{n-1} \|\Delta_k f\|_{L^p}. \end{aligned} \quad (2.5)$$

Now, Lemma 2.3 follows from Proposition 3.1 in [16], (2.4) and (2.5). $\square$

Lemma 2.4. [5] Let $1 \leq p \leq \infty$. Then for any $j \in \mathbb{Z}$, there hold

$$\begin{aligned} \|\Delta(\Delta_j f \circ \psi_j) - (\Delta \Delta_j f) \circ \psi_j\|_{L^p} &\leq C 2^{2j} e^{CV(t)} V(t) \|\Delta_j f\|_{L^p}, \\ \|\Lambda(\Delta_j f \circ \psi_j) - (\Lambda \Delta_j f) \circ \psi_j\|_{L^p} &\leq C 2^j e^{CV(t)} V(t)^{\frac{1}{2}} \|\Delta_j f\|_{L^p}. \end{aligned}$$

Here C is a constant independent of j, k .

2.2. The hybrid-Besov space. We denote the space $\mathcal{Z}'(\mathbb{R}^n)$ by the dual space of $\mathcal{Z}(\mathbb{R}^n) = \{f \in \mathcal{S}(\mathbb{R}^n); D^\alpha \hat{f}(0) = 0; \forall \alpha \in (\mathbb{N} \cup 0)^n \text{ multi-index}\}$. Let us first recall the definition of general Besov space.

Definition 2.5. Let $s \in \mathbb{R}$, $1 \leq p, q \leq +\infty$. The homogeneous Besov space $\dot{B}_{p,q}^s$ is defined by

$$\dot{B}_{p,q}^s \stackrel{\text{def}}{=} \{f \in \mathcal{Z}'(\mathbb{R}^n) : \|f\|_{\dot{B}_{p,q}^s} < +\infty\},$$

where

$$\|f\|_{\dot{B}_{p,q}^s} \stackrel{\text{def}}{=} \left\| 2^{ks} \|\Delta_k f(t)\|_{L^p} \right\|_{\ell^q}.$$

Now we introduce the hybrid-Besov space we will work with in this paper. Let $R_0 > 0$ be as in Proposition 4.4.

Definition 2.6. Let $s, \sigma \in \mathbb{R}$, $1 \leq p \leq +\infty$. The hybrid-Besov space $\dot{\mathcal{B}}_{2,p}^{s,\sigma}$ is defined by

$$\dot{\mathcal{B}}_{2,p}^{s,\sigma} \stackrel{\text{def}}{=} \{f \in \mathcal{Z}'(\mathbb{R}^n) : \|f\|_{\dot{\mathcal{B}}_{2,p}^{s,\sigma}} < +\infty\},$$

where

$$\|f\|_{\dot{\mathcal{B}}_{2,p}^{s,\sigma}} \stackrel{\text{def}}{=} \sum_{2^k \leq R_0} 2^{ks} \|\Delta_k f\|_{L^2} + \sum_{2^k > R_0} 2^{k\sigma} \|\Delta_k f\|_{L^p}.$$

We also denote $\dot{\mathcal{B}}^{s,\sigma}$ by $\dot{\mathcal{B}}_{2,2}^{s,\sigma}$.

The norm of the space $\tilde{L}_T^r(\dot{\mathcal{B}}_{2,p}^{s,\sigma})$ is defined by

$$\|f\|_{\tilde{L}_T^r(\dot{\mathcal{B}}_{2,p}^{s,\sigma})} \stackrel{\text{def}}{=} \sum_{2^k \leq R_0} 2^{ks} \|\Delta_k f\|_{L_T^r L^2} + \sum_{2^k > R_0} 2^{k\sigma} \|\Delta_k f\|_{L_T^r L^p}.$$

It is easy to check that $\tilde{L}_T^1(\dot{\mathcal{B}}_{2,p}^{s,\sigma}) = L_T^1(\dot{\mathcal{B}}_{2,p}^{s,\sigma})$ and $\tilde{L}_T^r(\dot{\mathcal{B}}_{2,p}^{s,\sigma}) \subseteq L_T^r(\dot{\mathcal{B}}_{2,p}^{s,\sigma})$ for $r > 1$.

The following lemma is a direct consequence of the definition of Besov space and Lemma 2.1.

Lemma 2.7. The following properties hold:

- (a) $\dot{\mathcal{B}}_{2,p}^{s_2,\sigma} \subseteq \dot{\mathcal{B}}_{2,p}^{s_1,\sigma}$ if $s_1 \geq s_2$ and $\dot{\mathcal{B}}_{2,p}^{s,\sigma_2} \subseteq \dot{\mathcal{B}}_{2,p}^{s,\sigma_1}$ if $\sigma_1 \leq \sigma_2$;
- (b) interpolation: for $s_1, s_2, \sigma_1, \sigma_2 \in \mathbb{R}$ and $\theta \in [0, 1]$, we have

$$\|f\|_{\dot{\mathcal{B}}_{2,p}^{\theta s_1 + (1-\theta)s_2, \theta \sigma_1 + (1-\theta)\sigma_2}} \leq \|f\|_{\dot{\mathcal{B}}_{2,p}^{s_1, \sigma_1}}^\theta \|f\|_{\dot{\mathcal{B}}_{2,p}^{s_2, \sigma_2}}^{(1-\theta)};$$

- (c) embedding: $\|f\|_{L^\infty} \leq C \|f\|_{\dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}}$;

- (d) inclusion relation: $\dot{\mathcal{B}}_{2,1}^s \subseteq \dot{\mathcal{B}}_{2,p}^{s, s - \frac{n}{2} + \frac{n}{p}} \subseteq \dot{\mathcal{B}}_{p,1}^{s - \frac{n}{2} + \frac{n}{p}}$ for $p \geq 2$.

Lemma 2.8. [14] Let $1 \leq p \leq \infty$. Then there hold

- (a) if $s_1, s_2 \leq \frac{n}{p}$ and $s_1 + s_2 > n \max(0, \frac{2}{p} - 1)$, then

$$\|fg\|_{\dot{\mathcal{B}}_{p,1}^{s_1+s_2-\frac{n}{p}}} \leq C \|f\|_{\dot{\mathcal{B}}_{p,1}^{s_1}} \|g\|_{\dot{\mathcal{B}}_{p,1}^{s_2}}.$$

- (b) if $s_1 \leq \frac{n}{p}, s_2 < \frac{n}{p}$, and $s_1 + s_2 \geq n \max(0, \frac{2}{p} - 1)$, then

$$\|fg\|_{\dot{\mathcal{B}}_{p,\infty}^{s_1+s_2-\frac{n}{p}}} \leq C \|f\|_{\dot{\mathcal{B}}_{p,1}^{s_1}} \|g\|_{\dot{\mathcal{B}}_{p,\infty}^{s_2}}.$$

Proposition 2.9. Let $\phi \in \mathcal{S}(\mathbb{R}^n)$, $p > n$. If $\phi_\varepsilon(x) \stackrel{\text{def}}{=} e^{i\frac{x_1}{\varepsilon}} \phi(x)$, then for any $\varepsilon > 0$,

$$\|\phi_\varepsilon\|_{\dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \leq C \varepsilon^{1-\frac{n}{p}},$$

here C is a constant independent of ε .

Proof. Fix $j_0 \in \mathbb{N}$ to be chosen later. By Lemma 2.1, we have

$$\sum_{j \geq j_0} 2^{(\frac{n}{p}-1)j} \|\Delta_j \phi_\varepsilon\|_{L^p} \leq C 2^{(\frac{n}{p}-1)j_0}.$$

Noting that $e^{i\frac{x_1}{\varepsilon}} = (-i\varepsilon\partial_1)^N e^{i\frac{x_1}{\varepsilon}}$ for any $N \in \mathbb{N}$, we get by integration by parts that

$$\Delta_j \phi_\varepsilon(x) = (i\varepsilon)^N 2^{jn} \int_{\mathbb{R}^n} e^{i\frac{y_1}{\varepsilon}} \partial_{y_1}^N (h(2^j(x-y))\phi(y)) dy, \quad h(x) \stackrel{\text{def}}{=} (\mathcal{F}^{-1}\varphi)(x),$$

from which and Young's inequality, we infer that

$$\|\Delta_j \phi_\varepsilon\|_{L^q} \leq C\varepsilon^N \max(2^{Nj}, 2^{(1-\frac{1}{q})nj}), \quad 1 \leq q \leq \infty,$$

which implies that by choosing $N > \frac{n}{p'}$

$$\begin{aligned} \sum_{0 \leq j < j_0} 2^{(\frac{n}{p}-1)j} \|\Delta_j \phi_\varepsilon\|_{L^p} &\leq C\varepsilon^N 2^{(N-1+\frac{n}{p})j_0}, \\ \sum_{j \leq \ln R_0} 2^{(\frac{n}{2}-1)j} \|\Delta_j \phi_\varepsilon\|_{L^2} &\leq C\varepsilon^N. \end{aligned}$$

Taking j_0 such that $2^{j_0} \sim \varepsilon^{-1}$ gives

$$\|\phi_\varepsilon\|_{\dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \leq C\varepsilon^{1-\frac{n}{p}}.$$

This completes the proof of Proposition 2.9. $\square$

3. REGULARIZING EFFECT OF THE HEAT EQUATION

Let us recall the classical parabolic regularizing effect.

Lemma 3.1. [7] Let ϕ be a smooth function supported in the annulus $\{\xi \in \mathbb{R}^n : A_1 \leq |\xi| \leq A_2\}$. Then there exist two positive constants c and C such that for any $1 \leq p \leq \infty$ and $\lambda > 0$, we have

$$\|\phi(\lambda^{-1}D)e^{\nu t\Delta}f\|_{L^p} \leq Ce^{-ct\lambda^2} \|\phi(\lambda^{-1}D)f\|_{L^p}.$$

Proposition 3.2. Let $s, \sigma \in \mathbb{R}$, and $p, r \in [1, \infty]$. Assume that $u_0 \in \dot{\mathcal{B}}_{2,p}^{s,\sigma}, f \in L_T^1 \dot{\mathcal{B}}_{2,p}^{s,\sigma}$. Then the heat equation

$$\begin{cases} \partial_t u - \nu \Delta u = f, \\ u|_{t=0} = u_0, \end{cases}$$

has a unique solution u satisfying

$$\|u\|_{\tilde{L}_T^r \dot{\mathcal{B}}_{2,p}^{s+\frac{2}{r}, \sigma+\frac{2}{r}}} \leq C(\|u_0\|_{\dot{\mathcal{B}}_{2,p}^{s,\sigma}} + \|f\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{s,\sigma}}).$$

Proof. The solution u can be written as

$$u(t) = e^{\nu t\Delta}u_0 + \int_0^t e^{\nu(t-s)\Delta}f(s)ds,$$

from which and Lemma 3.1, it follows that for any $1 \leq q \leq \infty$,

$$\|\Delta_j u(t)\|_{L^q} \leq Ce^{-ct2^{2j}} \|\Delta_j u_0\|_{L^q} + C \int_0^t e^{-c(t-s)2^{2j}} \|\Delta_j f(s)\|_{L^q} ds.$$

Taking L^r norm with respect to t gives

$$\|\Delta_j u(t)\|_{L_T^r L^q} \leq C2^{-\frac{2}{r}j} \|\Delta_j u_0\|_{L^q} + C2^{-\frac{2}{r}j} \|\Delta_j f(s)\|_{L_T^1 L^q}.$$

Using $q = 2$ for $2^j \leq R_0$ and $q = p$ for $2^j > R_0$, then Lemma follows from the definition of the hybrid-Besov space. $\square$

To prove the uniqueness of the solution, we need the following two propositions.

Proposition 3.3. [12] Let $p, q, r \in [1, \infty], s \in \mathbb{R}$. Assume that $u_0 \in \dot{B}_{p,q}^{s-1}, f \in L_T^1(\dot{B}_{p,q}^{s-1})$. Let u be a solution of the following equation

$$\partial_t u - \bar{\mu} \Delta u + (\bar{\lambda} + \bar{\mu}) \nabla \operatorname{div} u = f, \quad u(0, x) = u_0(x),$$

where $\bar{\mu} > 0, \bar{\lambda} + \bar{\mu} > 0$. Then for $t \in [0, T]$, there holds

$$\|u\|_{\tilde{L}_t^r(\dot{B}_{p,q}^{s-1+2/q})} \leq C(\|u_0\|_{\dot{B}_{p,q}^{s-1}} + \|f(\tau)\|_{\tilde{L}_t^1(\dot{B}_{p,q}^{s-1})}).$$

Proposition 3.4. [14] Let $s \in (-n \min(\frac{1}{p}, \frac{1}{p'}), 1 + \frac{n}{p})$, and $1 \leq p, q \leq +\infty$. Let v be a vector field such that $\nabla v \in L_T^1 \dot{B}_{p,1}^{\frac{n}{p}}$. Assume that $f_0 \in \dot{B}_{p,q}^s, g \in L_T^1(\dot{B}_{p,q}^s)$ and f is a solution of the transport equation

$$\partial_t f + v \cdot \nabla f = g, \quad f(0, x) = f_0.$$

Then for $t \in [0, T]$, there holds

$$\|f\|_{\tilde{L}_t^\infty(\dot{B}_{p,q}^s)} \leq e^{C \int_0^t \|\nabla v(\tau)\|_{\dot{B}_{p,1}^{\frac{n}{p}}} d\tau} \left(\|f_0\|_{\dot{B}_{p,q}^s} + \int_0^t \|g(\tau)\|_{\dot{B}_{p,q}^s} d\tau \right).$$

4. GREEN'S MATRIX OF THE LINEARIZED SYSTEM

We study the linearized system of the compressible Navier-Stokes system

$$\begin{cases} \partial_t a + \Lambda d = 0, \\ \partial_t d - \bar{\nu} \Delta d - \Lambda a = 0, \\ (a, d)|_{t=0} = (a_0, d_0). \end{cases} \quad (4.1)$$

We firstly give the explicit expression of Green's matrix for the system.

Lemma 4.1. Let $\mathcal{G}$ be the Green matrix of the system (4.1). Then we have the following explicit expression for $\hat{\mathcal{G}}$:

$$\hat{\mathcal{G}}(\xi, t) = \begin{bmatrix} \frac{\lambda_+ e^{\lambda_- t} - \lambda_- e^{\lambda_+ t}}{\lambda_+ - \lambda_-} & - \left(\frac{e^{\lambda_+ t} - e^{\lambda_- t}}{\lambda_+ - \lambda_-} \right) |\xi| \\ - \left(\frac{e^{\lambda_+ t} - e^{\lambda_- t}}{\lambda_+ - \lambda_-} \right) |\xi| & \frac{\lambda_+ e^{\lambda_+ t} - \lambda_- e^{\lambda_- t}}{\lambda_+ - \lambda_-} \end{bmatrix}, \quad (4.2)$$

where

$$\lambda_{\pm} = -\frac{1}{2} \bar{\nu} |\xi|^2 \pm \frac{1}{2} \sqrt{\bar{\nu}^2 |\xi|^4 - 4 |\xi|^2}.$$

Proof. We follow the proof of Lemma 3.1 in [20]. Firstly, applying the operator Λ to the second equation of (4.1) gives

$$(\Lambda d)_t + \Delta a = \bar{\nu} \Delta (\Lambda d).$$

Combining it with the first equation of (4.1), we get

$$a_{tt} = -(\Lambda d)_t = \Delta a + \bar{\nu} \Delta a_t.$$

Taking Fourier transform to the above equation yields that

$$\begin{cases} \hat{a}_{tt} + \bar{\nu} |\xi|^2 \hat{a}_t + |\xi|^2 \hat{a} = 0, \\ \hat{a}(\xi, 0) = \hat{a}_0(\xi), \quad \hat{a}_t(\xi, 0) = -|\xi| \hat{d}_0(\xi). \end{cases} \quad (4.3)$$

It is easy to check that $\lambda_{\pm}$ are two roots of the corresponding indicial equation of (4.3). Thus, we may assume that the solution of (4.3) has the form

$$\hat{a}(\xi, t) = A(\xi)e^{\lambda_{-}(\xi)t} + B(\xi)e^{\lambda_{+}(\xi)t}.$$

Using the initial conditions, we obtain

$$A = \frac{\lambda_{+}\hat{a}_0 + |\xi|\hat{d}_0}{\lambda_{+} - \lambda_{-}} \quad \text{and} \quad B = \frac{-|\xi|\hat{d}_0 - \lambda_{-}\hat{a}_0}{\lambda_{+} - \lambda_{-}},$$

which imply

$$\hat{a}(\xi, t) = \left(\frac{\lambda_{+}e^{\lambda_{-}t} - \lambda_{-}e^{\lambda_{+}t}}{\lambda_{+} - \lambda_{-}} \right) \hat{a}_0(\xi) - \left(\frac{e^{\lambda_{+}t} - e^{\lambda_{-}t}}{\lambda_{+} - \lambda_{-}} \right) |\xi| \hat{d}_0(\xi). \quad (4.4)$$

This determines $\widehat{\mathcal{G}}^{11}$ and $\widehat{\mathcal{G}}^{12}$.

On the other hand, taking Fourier transform to the second equation of (4.1) gives

$$\hat{d}_t = -\bar{\nu}|\xi|^2 \hat{d} + |\xi| \hat{a}.$$

Thus, we have

$$\hat{d}(\xi, t) = e^{-\bar{\nu}|\xi|^2 t} \left[\hat{d}(\xi, 0) + |\xi| \int_0^t e^{\bar{\nu}|\xi|^2 \tau} \hat{a}(\xi, \tau) d\tau \right].$$

Plugging (4.4) into the above equality and using the relations

$$\lambda_{\pm} + \bar{\nu}|\xi|^2 = -\lambda_{\mp}, \quad \lambda_{-}\lambda_{+} = |\xi|^2,$$

we finally get

$$\hat{d}(\xi, t) = - \left(\frac{e^{\lambda_{+}t} - e^{\lambda_{-}t}}{\lambda_{+} - \lambda_{-}} \right) |\xi| \hat{a}_0(\xi) + \left(\frac{\lambda_{+}e^{\lambda_{+}t} - \lambda_{-}e^{\lambda_{-}t}}{\lambda_{+} - \lambda_{-}} \right) \hat{d}_0(\xi),$$

which determines $\widehat{\mathcal{G}}^{21}$ and $\widehat{\mathcal{G}}^{22}$. □

We have the following pointwise estimates and expansion for $\widehat{\mathcal{G}}$.

Lemma 4.2. (a) Given $R > 0$, there is a positive number ϑ depending on R such that, for any multi-indices α and $|\xi| \leq R$,

$$|D_{\xi}^{\alpha} \widehat{\mathcal{G}}(\xi, t)| \leq C e^{-\vartheta|\xi|^2 t} (1 + |\xi|)^{|\alpha|} (1 + t)^{|\alpha|}, \quad (4.5)$$

where $C = C(R, |\alpha|)$;

(b) There exist positive constants R, ϑ depending on $\bar{\nu}$ such that the following expansion is valid for $|\xi| \geq R$,

$$\begin{aligned} \widehat{\mathcal{G}}(\xi, t) = & e^{-\bar{\nu}^{-1}t} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + e^{-\bar{\nu}|\xi|^2 t} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ & + \widehat{\mathcal{G}}^1(\xi, t) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \widehat{\mathcal{G}}^2(\xi, t) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \end{aligned} \quad (4.6)$$

where $\widehat{\mathcal{G}}^1$ and $\widehat{\mathcal{G}}^2$ satisfy the estimates

$$|\partial_{\xi}^{\alpha} \widehat{\mathcal{G}}^1| \leq C |\xi|^{-|\alpha|-1} (e^{-\frac{\bar{\nu}^{-1}t}{2}} + e^{-\vartheta|\xi|^2 t}), \quad (4.7)$$

$$|\partial_{\xi}^{\alpha} \widehat{\mathcal{G}}^2| \leq C |\xi|^{-|\alpha|-2} (e^{-\frac{\bar{\nu}^{-1}t}{2}} + e^{-\vartheta|\xi|^2 t}), \quad (4.8)$$

for a positive constant C depending on $|\alpha|, \bar{\nu}$.

Remark 4.3. In fact, $\widehat{\mathcal{G}}^2$ is a diagonal matrix, see (4.9). Since both nonzero elements in $\widehat{\mathcal{G}}^2$ can be estimated by the right side of (4.8), we don't care about its explicit expression, and we view it as a scalar function.

Proof. We follow the proof of Lemma 3.2 in [20]. Since the explicit expansion of $\widehat{\mathcal{G}}$ is important for us, here we will present a proof. We refer to [20] for more details.

The inequality (4.5) can be deduced from Lemma 4.1 by routine calculations. We present the proof of (b) below. Let p be the symbol

$$p(z, r) = z^2 + \bar{\nu}r^2z + r^2.$$

Define contours Γ_+, Γ_- and Γ_0 in the complex plane to be the circles of radius $\frac{\bar{\nu}^{-1}}{2}$ centered respectively at $-\bar{\nu}^{-1}$, $-\bar{\nu}r^2 + \bar{\nu}^{-1}$ and 0, and

$$\begin{cases} E(r, t) = \frac{1}{2\pi i} \int_{\Gamma_- \cup \Gamma_+} \frac{e^{tz}}{p(z, r)} dz, \\ F(r, t) = E_t(r, t) + \bar{\nu}r^2 E(r, t), \end{cases}$$

for large enough such that $r > 10\bar{\nu}^{-1}$. Then it is easy to verify that

$$\begin{cases} p(\partial_t, r)E = p(\partial_t, r)F = 0, \\ E(r, 0) = 0, \quad F(r, 0) = 1, \\ E_t(r, 0) = 1, \quad F_t(r, 0) = 0. \end{cases}$$

Thus, we may assume that E and F have the form

$$\begin{aligned} E(\xi, t) &= A_E(\xi)e^{\lambda_- t} + B_E(\xi)e^{\lambda_+ t}, \\ F(\xi, t) &= A_F(\xi)e^{\lambda_- t} + B_F(\xi)e^{\lambda_+ t}. \end{aligned}$$

Thanks to the initial conditions on E and F , we obtain

$$A_E = \frac{-1}{\lambda_+ - \lambda_-}, \quad B_E = \frac{1}{\lambda_+ - \lambda_-}, \quad A_F = \frac{\lambda_+}{\lambda_+ - \lambda_-}, \quad B_F = \frac{-\lambda_-}{\lambda_+ - \lambda_-}.$$

Let $\hat{a}, \hat{d}$ be as in the proof of Lemma 4.1. By (4.3), we find

$$\hat{a}(\xi, t) = F(|\xi|, t)\hat{a}_0(\xi) - |\xi|E(|\xi|, t)\hat{d}_0(\xi), \quad \widehat{\mathcal{G}}^{21} = \widehat{\mathcal{G}}^{12},$$

which implies that

$$\widehat{\mathcal{G}}^{11}(\xi, t) = F(|\xi|, t), \quad \widehat{\mathcal{G}}^{12}(\xi, t) = -|\xi|E(|\xi|, t), \quad \widehat{\mathcal{G}}^{21}(\xi, t) = -|\xi|E(|\xi|, t).$$

Introducing the function

$$H(r, t) = e^{-\bar{\nu}r^2 t} \int_0^t e^{\bar{\nu}r^2 \tau} E(r, \tau) d\tau = \frac{\lambda_- e^{\lambda_- t} - \lambda_+ e^{\lambda_+ t}}{|\xi|^2(\lambda_+ - \lambda_-)},$$

then we find

$$\widehat{\mathcal{G}}^{22}(\xi, t) = e^{-\bar{\nu}|\xi|^2 t} - |\xi|^2 H(\xi, t).$$

Thus we have

$$\begin{aligned} \widehat{\mathcal{G}} &= \begin{bmatrix} e^{-\bar{\nu}^{-1}t} & 0 \\ 0 & e^{-\bar{\nu}|\xi|^2 t} \end{bmatrix} - |\xi|E(|\xi|, t) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &\quad + \begin{bmatrix} F(|\xi|, t) - e^{-\bar{\nu}^{-1}t} & 0 \\ 0 & -|\xi|^2 H(\xi, t) \end{bmatrix}. \end{aligned}$$

Consequently, we obtain the required expansion (4.6) of $\widehat{G}$ with

$$\widehat{\mathcal{G}}^1(\xi, t) = -|\xi|E(|\xi|, t), \quad \widehat{\mathcal{G}}^2(\xi, t) = \begin{bmatrix} F(|\xi|, t) - e^{-\bar{\nu}^{-1}t} & 0 \\ 0 & -|\xi|^2 H(\xi, t) \end{bmatrix}. \quad (4.9)$$

It remains to prove (4.7) and (4.8). By making a change $w = z + \bar{\nu}r^2 - \bar{\nu}^{-1}$, we get

$$\begin{aligned} r \int_{\Gamma_-(r)} \frac{e^{tz}}{p(z, r)} dz &= -e^{(\bar{\nu}^{-1} - \bar{\nu}r^2)t} \int_{\Gamma_0} \frac{e^{tw}}{\bar{\nu}rw} \left[1 - \frac{(w + \bar{\nu}^{-1})^2}{\bar{\nu}r^2w} \right]^{-1} dw \\ &= \sum_{j=0}^{\infty} r^{-2j-1} \left[-e^{(\bar{\nu}^{-1} - \bar{\nu}r^2)t} \int_{\Gamma_0} \frac{(w + \bar{\nu}^{-1})^{2j}}{(\bar{\nu}w)^{j+1}} e^{tw} dw \right]. \end{aligned}$$

for large enough $r > 10\bar{\nu}^{-1}$, here C is an absolute constant. Let $b_j(t)$ be the contour integral on the right hand side, we have

$$|b_j(t)| \leq C(\bar{\nu}) e^{\frac{\bar{\nu}^{-1}t}{2}} (\widetilde{C}\bar{\nu}^{-1})^{2j},$$

where $\widetilde{C}$ is an absolutely constant. Then if $r > 2\bar{\nu}^{-1}$ is large enough, the term in bracket is bounded by

$$C(\bar{\nu}) e^{(\bar{\nu}^{-1} - \bar{\nu}r^2)t} e^{\frac{\bar{\nu}^{-1}t}{2}} (\widetilde{C}\bar{\nu}^{-1})^{2j} \leq C(\bar{\nu}) e^{-\frac{\bar{\nu}r^2t}{2}} (\widetilde{C}\bar{\nu}^{-1})^{2j}.$$

Taking k -derivatives with respect to r , we have

$$\left(\frac{\partial}{\partial r} \right)^k \int_{\Gamma_-} \frac{re^{tz}}{p(z, r)} dz = -e^{\bar{\nu}^{-1}t} \sum_{j=0}^{\infty} b_j(t) \sum_{\ell=0}^k \binom{k}{\ell} \left[\left(\frac{\partial}{\partial r} \right)^{\ell} e^{-\bar{\nu}r^2t} \right] \left[\left(\frac{\partial}{\partial r} \right)^{k-\ell} r^{-2j-1} \right].$$

Noting that $\partial_r^{\ell} e^{-\bar{\nu}r^2t} \leq C(\ell) r^{-\ell} e^{-\frac{2}{3}\bar{\nu}r^2t}$, so if $r > 2\widetilde{C}\bar{\nu}^{-1}$ is large enough, we have

$$\begin{aligned} &\left| \left(\frac{\partial}{\partial r} \right)^k \int_{\Gamma_-} \frac{re^{tz}}{p(z, r)} dz \right| \\ &\leq C e^{(\bar{\nu}^{-1} - \frac{2}{3}\bar{\nu}r^2)t} \sum_{j=0}^{\infty} |b_j(t)| r^{-2j-1-k} \sum_{\ell=0}^k \binom{k}{\ell} \frac{(2j+k-\ell)!}{(2j)!} \\ &\leq C(\bar{\nu}, k) e^{-\frac{1}{2}\bar{\nu}r^2t} r^{-k-1} \sum_{j=0}^{\infty} \left(\frac{\widetilde{C}\bar{\nu}^{-1}}{r} \right)^{2j} \frac{(2j+k-\ell)!}{(2j)!} \\ &\leq C(\bar{\nu}, k) e^{-\frac{1}{2}\bar{\nu}r^2t} r^{-k-1}. \end{aligned} \quad (4.10)$$

On the other hand, by making a change $w = z + \bar{\nu}^{-1}$, we have

$$\begin{aligned} r \int_{\Gamma_+(r)} \frac{e^{tz}}{p(z, r)} dz &= e^{-\bar{\nu}^{-1}t} \int_{\Gamma_0} \frac{e^{tw}}{\bar{\nu}r^2w} \left[1 - \frac{(w - \bar{\nu}^{-1})^2}{\bar{\nu}rw} \right]^{-1} dw \\ &= \sum_{j=0}^{\infty} r^{-2j-1} (-1)^j \left[e^{-\bar{\nu}^{-1}t} \int_{\Gamma_0} \frac{(w - \bar{\nu}^{-1})^{2j}}{(\bar{\nu}w)^{j+1}} e^{tw} dw \right]. \end{aligned}$$

The term in the bracket is dominated by

$$C(\bar{\nu}) e^{-\frac{\bar{\nu}^{-1}t}{2}} (\widetilde{C}\bar{\nu}^{-1})^{2j},$$

then by the same argument as leading to (4.10), we get

$$\left| \left(\frac{\partial}{\partial r} \right)^k \int_{\Gamma_+} \frac{r e^{tz}}{p(z, r)} dz \right| \leq C(\bar{\nu}) e^{-\frac{\bar{\nu}-1}{2} t} r^{-k-1}. \quad (4.11)$$

With (4.10) and (4.11), we can obtain (4.7) and (4.8) by using the definitions of the functions E, F and H . $\square$

Using Lemma 4.2, we can obtain the following smoothing effect of Green's matrix $\mathcal{G}$, which will play an important role in this paper.

Proposition 4.4. Let $\mathcal{C}$ be a ring centered at 0 in $\mathbb{R}^n$. Then there exist positive constants R_0, C, c depending on $\bar{\nu}$ such that, if $\text{supp } \hat{u} \subset \lambda \mathcal{C}$, then we have

(a) if $\lambda \leq R_0$, then

$$\|\mathcal{G} * u\|_{L^2} \leq C e^{-c\lambda^2 t} \|u\|_{L^2}; \quad (4.12)$$

(b) if $b \leq \lambda \leq R_0$, then for any $1 \leq p \leq \infty$,

$$\|\mathcal{G} * u\|_{L^p} \leq C(1 + b^{-n-1}) e^{-c\lambda^2 t} \|u\|_{L^p}; \quad (4.13)$$

(c) if $\lambda > R_0$, then for any $1 \leq p \leq \infty$,

$$\|\mathcal{G}^1 * u\|_{L^p} \leq C \lambda^{-1} e^{-ct} \|u\|_{L^p}, \quad (4.14)$$

$$\|\mathcal{G}^2 * u\|_{L^p} \leq C \lambda^{-2} e^{-ct} \|u\|_{L^p}. \quad (4.15)$$

Proof. (a) Thanks to Plancherel theorem, we get

$$\|\mathcal{G} * u\|_{L^2} = \|\widehat{\mathcal{G}}(\xi) \hat{u}(\xi)\|_{L^2} \leq C \|e^{-\vartheta|\xi|^2 t} \hat{u}(\xi)\|_2 \leq C e^{-c\lambda^2 t} \|u\|_2,$$

where we have used (4.5) and the support property of $\hat{u}(\xi)$.

(b) Let $\phi \in \mathcal{D}(\mathbb{R}^n \setminus \{0\})$, which equals to 1 near the ring $\mathcal{C}$. Set

$$g(t, x) \stackrel{\text{def}}{=} (2\pi)^{-n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \phi(\lambda^{-1} \xi) \widehat{\mathcal{G}}(\xi, t) d\xi.$$

To prove (4.13), it suffices to show

$$\|g(x, t)\|_{L^1} \leq C(1 + b^{-n-1}) e^{-c\lambda^2 t}. \quad (4.16)$$

Thanks to (4.5), we infer that

$$\int_{|x| \leq \lambda^{-1}} |g(x, t)| dx \leq C \int_{|x| \leq \lambda^{-1}} \int_{\mathbb{R}^n} |\phi(\lambda^{-1} \xi)| |\widehat{\mathcal{G}}(\xi, t)| d\xi dx \leq C e^{-c\lambda^2 t}. \quad (4.17)$$

Set $L \stackrel{\text{def}}{=} \frac{x \cdot \nabla_\xi}{i|x|^2}$. Noting that $L(e^{ix \cdot \xi}) = e^{ix \cdot \xi}$, we get by integration by part that

$$\begin{aligned} g(x, t) &= \int_{\mathbb{R}^n} L^{n+1}(e^{ix \cdot \xi}) \phi(\lambda^{-1} \xi) \widehat{\mathcal{G}}(\xi, t) d\xi \\ &= (-1)^{n+1} \int_{\mathbb{R}^n} e^{ix \cdot \xi} (L^*)^{n+1}(\phi(\lambda^{-1} \xi) \widehat{\mathcal{G}}(\xi, t)) d\xi. \end{aligned}$$

From the Leibnitz's formula and (4.5),

$$\begin{aligned} & |(L^*)^{n+1}(\phi(\lambda^{-1} \xi) \widehat{\mathcal{G}}(\xi, t))| \\ & \leq C |\lambda x|^{-(n+1)} \sum_{|\gamma|=n+1, |\beta| \leq |\gamma|} \lambda^{|\beta|} |(\nabla^{\gamma-\beta} \phi)(\lambda^{-1} \xi)| e^{-\vartheta|\xi|^2 t} (1 + |\xi|)^{|\beta|} (1 + t)^{|\beta|}. \end{aligned}$$

Thanks to the estimate

$$\begin{aligned} e^{-\vartheta|\xi|^2 t}(1+|\xi|)^{|\beta|}(1+t)^{|\beta|} &\leq C e^{-\vartheta|\xi|^2 t}(1+t^{|\beta|}+|\xi|^{|\beta|}+t^{|\beta|}|\xi|^{|\beta|}) \\ &\leq C e^{-\frac{\vartheta}{2}|\xi|^2 t}(1+|\xi|^{-2|\beta|}+|\xi|^{|\beta|}+|\xi|^{-|\beta|}), \end{aligned}$$

we obtain, for any ξ with $b \lesssim |\xi| \sim \lambda \leq R_0$,

$$|(L^*)^{n+1}(\phi(\lambda^{-1}\xi)\widehat{\mathcal{G}}(\xi, t))| \leq C(1+b^{-n-1})|\lambda x|^{-(n+1)}e^{-\frac{\vartheta}{2}|\xi|^2 t},$$

which implies that

$$\begin{aligned} \int_{|x| \geq \frac{1}{\lambda}} |g(x, t)| dx &\leq C(1+b^{-n-1})e^{-c\lambda^2 t} \lambda^n \int_{|x| \geq \frac{1}{\lambda}} |\lambda x|^{-n-1} dx \\ &\leq C(1+b^{-n-1})e^{-c\lambda^2 t}, \end{aligned}$$

which together with (4.17) gives (4.16).

(c) We set

$$g^1(t, x) \stackrel{\text{def}}{=} (2\pi)^{-n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \phi(\lambda^{-1}\xi) \widehat{\mathcal{G}}^1(\xi, t) d\xi.$$

Since the integrand is supported in $\{\xi : |\xi| \sim \lambda > R_0\}$, then we get by (4.7) that

$$\begin{aligned} \int_{|x| \leq \lambda^{-1}} |g^1| dx &\leq C \int_{|x| \leq \lambda^{-1}} \int |\phi(\lambda^{-1}\xi)| |\xi|^{-1} (e^{-\frac{\bar{\nu}-1}{2}t} + e^{-\vartheta|\xi|^2 t}) d\xi dx \\ &\leq C \lambda^{-1} e^{-ct}. \end{aligned} \tag{4.18}$$

We get by integration by parts that

$$g^1(x, t) = (-1)^{n+1} \int_{\mathbb{R}^n} e^{ix \cdot \xi} (L^*)^{n+1}(\phi(\lambda^{-1}\xi) \widehat{\mathcal{G}}^1(\xi, t)) d\xi.$$

While thanks to (4.7), the integrand is dominated by

$$|\lambda x|^{-n-1} \lambda^{-1} e^{-ct},$$

which implies that

$$\begin{aligned} \int_{|x| \geq \lambda^{-1}} |g^1(x, t)| dx &\leq C \lambda^{-1} e^{-ct} \lambda^n \int_{|x| \geq \lambda^{-1}} |\lambda x|^{-n-1} dx \\ &\leq C \lambda^{-1} e^{-ct}, \end{aligned} \tag{4.19}$$

from which and (4.18), it follows that

$$\|g^1(x, t)\|_{L^1} \leq C \lambda^{-1} e^{-ct},$$

which implies (4.14). The inequality (4.15) can be similarly proved. $\square$

5. THE LINEARIZED SYSTEM WITH CONVECTION TERM

In this section we consider the linearized system with convection terms

$$\begin{cases} \partial_t a + \Lambda d = -v \cdot \nabla a + F, \\ \partial_t d - \bar{\nu} \Delta d - \Lambda a = -v \cdot \nabla d + G, \\ (a, d)|_{t=0} = (a_0, d_0). \end{cases} \tag{5.1}$$

Let $T > 0$, $s \in \mathbb{R}$, $p \geq 2$, and $s_p \stackrel{\text{def}}{=} s - \frac{n}{p} + \frac{n}{2}$. We introduce the functional space $\mathcal{E}_T^s$ which is defined as follows

$$\mathcal{E}_T^s \stackrel{\text{def}}{=} \left\{ (a, d) \in \left(L^1(0, T; \dot{\mathcal{B}}_{2,p}^{s_p+1,s}) \cap \tilde{L}^\infty(0, T; \dot{\mathcal{B}}_{2,p}^{s_p-1,s}) \right) \right. \\ \left. \times \left(L^1(0, T; \dot{\mathcal{B}}_{2,p}^{s_p+1,s+1}) \cap \tilde{L}^\infty(0, T; \dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}) \right)^n \right\}$$

with the norm

$$\|(a, d)\|_{\mathcal{E}_T^s} \stackrel{\text{def}}{=} \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{s_p-1,s}} + \|d\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}} + \|a\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p+1,s}} + \|d\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p+1,s+1}}.$$

We denote

$$\overline{V}(t) \stackrel{\text{def}}{=} \|v\|_{L_t^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} + \|v\|_{\tilde{L}_t^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}.$$

Theorem 5.1. Let $2 \leq p < 2n$, $p \leq \min(4, \frac{2n}{n-2})$, and $1 - \frac{n}{p} < s \leq 1 + \frac{2n}{p} - \frac{n}{2}$. Assume that $v \in \tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1} \cap L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}$, $F \in L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p-1,s}$, $G \in L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}$. Let (a, d) be a solution of (5.1) on $[0, T]$. Then there exists a constant C independent of T such that

$$\|(a, d)\|_{\mathcal{E}_T^s} \leq C e^{C\overline{V}(T)} \left\{ \|(a_0, d_0)\|_{\mathcal{E}_0^s} + (\overline{V}(T) + \overline{V}(T)^{\frac{1}{2}}) \|(a, d)\|_{\mathcal{E}_T^s} \right. \\ \left. + \|F\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p-1,s}} + \|G\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}} \right\}.$$

Here $\|(a_0, d_0)\|_{\mathcal{E}_0^s} \stackrel{\text{def}}{=} \|a_0\|_{\dot{\mathcal{B}}_{2,p}^{s_p-1,s}} + \|d_0\|_{\dot{\mathcal{B}}_{2,p}^{s_p-1,s-1}}$.

We introduce another functional space E_T^s which is defined by

$$E_T^s \stackrel{\text{def}}{=} \left\{ (a, d) \in \left(L^1(0, T; \dot{\mathcal{B}}^{s+1,s}) \cap \tilde{L}^\infty(0, T; \dot{\mathcal{B}}^{s-1,s}) \right) \right. \\ \left. \times \left(L^1(0, T; \dot{\mathcal{B}}_{2,1}^{s+1}) \cap \tilde{L}^\infty(0, T; \dot{\mathcal{B}}_{2,1}^{s-1}) \right)^n \right\}$$

with the norm

$$\|(a, d)\|_{E_T^s} \stackrel{\text{def}}{=} \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}^{s-1,s}} + \|d\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,1}^{s-1}} + \|a\|_{L_T^1 \dot{\mathcal{B}}^{s+1,s}} + \|d\|_{L_T^1 \dot{\mathcal{B}}_{2,1}^{s+1}}.$$

Theorem 5.2. Let $2 \leq p < 2n$, $p \leq \min(4, \frac{2n}{n-2})$, and $1 - \frac{n}{p} < s \leq 1 + \frac{n}{p}$. Assume that $v \in \tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1} \cap L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}$, $F \in L_T^1 \dot{\mathcal{B}}^{s-1,s}$, $G \in L_T^1 \dot{\mathcal{B}}_{2,1}^{s-1}$. Let (a, d) be a solution of (5.1) on $[0, T]$. Then there exists a constant C independent of T such that

$$\|(a, d)\|_{E_T^s} \leq C e^{C\overline{V}(T)} \left\{ \|(a_0, d_0)\|_{E_0^s} + (\overline{V}(T) + \overline{V}(T)^{\frac{1}{2}}) \|(a, d)\|_{E_T^s} \right. \\ \left. + \|v\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,1}^s} \|(a, d)\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} + \|F\|_{L_T^1 \dot{\mathcal{B}}_{2,1}^{s-1,s}} + \|G\|_{L_T^1 \dot{\mathcal{B}}_{2,1}^{s-1}} \right\}.$$

Here $\|(a_0, d_0)\|_{E_0^s} \stackrel{\text{def}}{=} \|a_0\|_{\dot{\mathcal{B}}^{s-1,s}} + \|d_0\|_{\dot{\mathcal{B}}_{2,1}^{s-1}}$.

We denote

$$a_j \stackrel{\text{def}}{=} \Delta_j a, \quad d_j \stackrel{\text{def}}{=} \Delta_j d, \quad F_j \stackrel{\text{def}}{=} \Delta_j F, \quad G_j \stackrel{\text{def}}{=} \Delta_j G.$$

The proof of Theorem 5.1-5.2 is based on the following frequency localized system:

$$\begin{cases} \partial_t a_j + \Lambda d_j = -\Delta_j(v \cdot \nabla a) + F_j, \\ \partial_t d_j - \bar{\nu} \Delta d_j - \Lambda a_j = -\Delta_j(v \cdot \nabla d) + G_j, \\ (a_j, d_j)|_{t=0} = (\Delta_j a_0, \Delta_j d_0) \stackrel{\text{def}}{=} (a_j^0, d_j^0). \end{cases} \quad (5.2)$$

In what follows, we define $\tilde{p}$ as $\frac{1}{\tilde{p}} = \frac{n}{2p} - \frac{n}{4} + \frac{1}{2}$, and denote $p', \tilde{p}'$ by the conjugate index of $p, \tilde{p}$ respectively.

5.1. Low frequency estimate: $2^j \leq R_0$. In this case, the Green matrix of the linearized system behaves as the heat kernel. Using the smoothing effect of the Green matrix, the terms $v \cdot \nabla a$ and $v \cdot \nabla d$ can be handled as the perturbation terms.

In terms of the Green matrix, the solution of (5.2) can be expressed as

$$\begin{pmatrix} a_j \\ d_j \end{pmatrix} = \mathcal{G}(x, t) * \begin{pmatrix} a_j^0 \\ d_j^0 \end{pmatrix} + \int_0^t \mathcal{G}(x, t - \tau) * \begin{pmatrix} F_j - \Delta_j(v \cdot \nabla a) \\ G_j - \Delta_j(v \cdot \nabla d) \end{pmatrix} d\tau.$$

From Proposition 4.4 (a) and Young's inequality, we infer that

$$\begin{aligned} \|a_j(t)\|_{L^2} + \|d_j(t)\|_{L^2} &\leq C e^{-c2^{2j}t} (\|a_j^0\|_{L^2} + \|d_j^0\|_{L^2}) \\ &\quad + C \int_0^t e^{-c2^{2j}(t-\tau)} (\|F_j(\tau)\|_{L^2} + \|G_j(\tau)\|_{L^2}) d\tau \\ &\quad + C \int_0^t e^{-c2^{2j}(t-\tau)} (\|\Delta_j(v \cdot \nabla a)(\tau)\|_{L^2} + \|\Delta_j(v \cdot \nabla d)(\tau)\|_{L^2}) d\tau. \end{aligned}$$

Taking L^r norm with respect to t gives

$$\begin{aligned} \|a_j\|_{L_T^r L^2} + \|d_j\|_{L_T^r L^2} &\leq C 2^{-\frac{2j}{r}} (\|a_j^0\|_{L^2} + \|d_j^0\|_{L^2} + \|F_j\|_{L_T^1 L^2} + \|\Delta_j(v \cdot \nabla a)\|_{L_T^1 L^2} \\ &\quad + \|G_j\|_{L_T^1 L^2} + \|\Delta_j(v \cdot \nabla d)\|_{L_T^1 L^2}). \end{aligned}$$

Here and what follows, we assume $1 \leq r \leq \infty$. Noting that $1 - \frac{n}{p} < s \leq \frac{2n}{p} - \frac{n}{2} + 1$, we apply Proposition 7.3 (b) with $s = \frac{n}{p}, t = s - \frac{n}{p} + \frac{n}{p'} - 1, \tilde{s} = s_p - 1, \tilde{t} = \frac{n}{2}, \gamma = \frac{n}{p} - \frac{n}{2}$ to get

$$\begin{aligned} &\sum_{2^j \leq R_0} 2^{j(s_p-1)} \|\Delta_j(v \nabla a)\|_{L_T^1 L^2} \\ &\leq C \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \|\nabla a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s - \frac{n}{p} + \frac{n}{p'} - 1, s_p - 1 + \gamma}} + C \|\nabla a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{s_p-1, s-1}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \\ &\leq C \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \|a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s - \frac{n}{p} + \frac{n}{p'}, s}} + C \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{s_p, s}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}}, \end{aligned}$$

and with $\gamma = 0$,

$$\begin{aligned} &\sum_{2^j \leq R_0} 2^{j(s_p-1)} \|\Delta_j(v \nabla d)\|_{L_T^1 L^2} \\ &\leq C \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \|\nabla d\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s - \frac{n}{p} + \frac{n}{p'} - 1, s_p - 1 + \gamma}} + C \|\nabla d\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{s_p-1, s-1}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \\ &\leq C \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \|d\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s - \frac{n}{p} + \frac{n}{p'}, s_p}} + C \|d\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{s_p, s}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}}. \end{aligned}$$

Then we conclude that for any $1 \leq r \leq \infty$,

$$\begin{aligned}
& \sum_{2^j \leq R_0} 2^{j(s_p-1+\frac{2}{r})} (\|a_j\|_{L_T^r L^2} + \|d_j\|_{L_T^r L^2}) \\
& \leq C \sum_{2^j \leq R_0} 2^{j(s_p-1)} (\|a_j^0\|_{L^2} + \|d_j^0\|_{L^2}) \\
& \quad + C \left\{ \|F\|_{L_T^1 \dot{B}_{2,p}^{s_p-1,s}} + \|G\|_{L_T^1 \dot{B}_{2,p}^{s_p-1,s-1}} + \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2},\frac{n}{p}}} \|(a,d)\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{s_p,s}} \right. \\
& \quad \left. + \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p},\frac{2n}{p}-\frac{n}{2}}} \left(\|a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p}+\frac{n}{p'},s}} + \|d\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p}+\frac{n}{p'},s_p}} \right) \right\}. \quad (5.3)
\end{aligned}$$

5.2. High frequency estimate: $2^j > R_0$. In this case, since the Green's matrix has no smoothing effect on a , we will lose one derivative about a in the energy estimate if we still regard $v \cdot \nabla a$ as the perturbation term. To avoid the loss of the derivative, we will work in the Lagrangian coordinate in this subsection.

Firstly, we rewrite (5.2) as

$$\begin{cases} \partial_t a_j + S_{j-1} v \cdot \nabla a_j + \Lambda d_j = (S_{j-1} v \cdot \nabla a_j - \Delta_j(v \cdot \nabla a)) + F_j, \\ \partial_t d_j + S_{j-1} v \cdot \nabla d_j - \bar{\nu} \Delta d_j - \Lambda a_j = (S_{j-1} v \cdot \nabla d_j - \Delta_j(v \cdot \nabla d)) + G_j, \\ (a_j, d_j)|_{t=0} = (a_j^0, d_j^0). \end{cases} \quad (5.4)$$

Let $\psi_j(t, x)$ be the solution to

$$\frac{d}{dt} \psi_j(t, x) = S_{j-1} v(t, \psi_j(t, x)), \quad \psi_j(0, x) = x.$$

We set

$$\bar{a}_j = a_j(t, \psi_j(t, x)), \quad \bar{d}_j = d_j(t, \psi_j(t, x)), \quad \bar{F}_j = F_j(t, \psi_j(t, x)), \quad \bar{G}_j = G_j(t, \psi_j(t, x)).$$

Then the new unknown $(\bar{a}_j, \bar{d}_j)$ satisfies

$$\begin{cases} \partial_t \bar{a}_j + \Lambda \bar{d}_j = \bar{F}_j + \bar{\mathcal{R}}_j, \\ \partial_t \bar{d}_j - \bar{\nu} \Delta \bar{d}_j - \Lambda \bar{a}_j = \bar{G}_j + \bar{\mathcal{Q}}_j, \\ (\bar{a}_j, \bar{d}_j)|_{t=0} = (\Delta_j a_0, \Delta_j d_0), \end{cases} \quad (5.5)$$

where

$$\begin{aligned}
\bar{\mathcal{R}}_j & \stackrel{\text{def}}{=} (S_{j-1} v \cdot \nabla a_j - \Delta_j(v \cdot \nabla a)) \circ \psi_j + \Lambda(d_j \circ \psi_j) - (\Lambda d_j) \circ \psi_j, \\
\bar{\mathcal{Q}}_j & \stackrel{\text{def}}{=} ((S_{j-1} v \cdot \nabla d_j) - \Delta_j(v \cdot \nabla d)) \circ \psi_j - \bar{\nu} \Delta(d_j \circ \psi_j) + \bar{\nu}(\Delta d_j) \circ \psi_j \\
& \quad + (\Lambda a_j) \circ \psi_j - \Lambda(a_j \circ \psi_j).
\end{aligned}$$

We get by Lemma 2.2 that

$$\|a_j\|_{L^p} \leq e^{\bar{V}(t)} \|\bar{a}_j\|_{L^p}, \quad (5.6)$$

where we used the fact

$$\int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau \leq C \int_0^t \|\nabla v(\tau)\|_{\dot{B}_{2,p}^{\frac{n}{2},\frac{n}{p}}} d\tau \leq C \bar{V}(t).$$

Fix $N \in \mathbb{N}$ to be chosen later. Noting that

$$\bar{a}_j = \sum_k \Delta_k \bar{a}_j = \sum_{|k-j| \leq N} \Delta_k \bar{a}_j + \sum_{|k-j| > N} \Delta_k \bar{a}_j + S_{j-N} \bar{a}_j,$$

we have by (5.6) that

$$\|a_j\|_{L^p} \leq e^{\bar{V}(t)} \left(\sum_{|k-j| \leq N} \|\Delta_k \bar{a}_j\|_{L^p} + \sum_{k-j > N} \|\Delta_k \bar{a}_j\|_{L^p} + \|S_{j-N} \bar{a}_j\|_{L^p} \right).$$

Thanks to Lemma 2.3, we have

$$\begin{aligned} \sum_{k-j > N} \|\Delta_k \bar{a}_j\|_{L^p} &\leq C e^{C\bar{V}(t)} \sum_{k-j > N} 2^{-(k-j)} \|a_j\|_{L^p} \leq C 2^{-N} e^{C\bar{V}(t)} \|a_j\|_{L^p}, \\ \|S_{j-N} \bar{a}_j\|_{L^p} &\leq C e^{C\bar{V}(t)} (\bar{V}(t) \|a_j\|_{L^p} + 2^{-N} \|a_j\|_{L^p}). \end{aligned}$$

Summing up the above estimates yields

$$\|a_j\|_{L^p} \leq C e^{C\bar{V}(t)} \left(\sum_{|k-j| \leq N} \|\Delta_k \bar{a}_j\|_{L^p} + 2^{-N} \|a_j\|_{L^p} + \bar{V}(t) \|a_j\|_{L^p} \right).$$

Choosing N large enough such that

$$C e^{C\bar{V}(T)} 2^{-N} \leq \frac{1}{2} \quad \text{i.e., } N \sim e^{C\bar{V}(T)},$$

then we obtain

$$\|a_j\|_{L_T^r L^p} \leq C e^{C\bar{V}(T)} \left(\sum_{|k-j| \leq N} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} + \bar{V}(T) \|a_j\|_{L_T^r L^p} \right). \quad (5.7)$$

Similarly, we can obtain

$$\|d_j\|_{L_T^r L^p} \leq C e^{C\bar{V}(T)} \left(\sum_{|k-j| \leq N} \|\Delta_k \bar{d}_j\|_{L_T^r L^p} + \bar{V}(T) \|d_j\|_{L_T^r L^p} \right). \quad (5.8)$$

Next, we need to estimate $\|\Delta_k \bar{a}_j\|_{L_T^r L^p}$ and $\|\Delta_k \bar{d}_j\|_{L_T^r L^p}$ for $|k-j| \leq N$. In order to do this, we first present the estimates of $\bar{F}_j, \bar{G}_j, \bar{\mathcal{R}}_j$ and $\bar{\mathcal{Q}}_j$. We get by using Lemma 2.3 that

$$\begin{aligned} \|\Delta_k \bar{F}_j\|_{L_T^1 L^p} &\leq C 2^{-(k-j)} e^{C\bar{V}(T)} \|F_j\|_{L_T^1 L^p}, \\ \|\Delta_k \bar{G}_j\|_{L_T^1 L^p} &\leq C 2^{-(k-j)} e^{C\bar{V}(T)} \|G_j\|_{L_T^1 L^p}. \end{aligned} \quad (5.9)$$

Set

$$\begin{aligned} \bar{\mathcal{R}}_j^1 &= (S_{j-1} v \cdot \nabla a_j - \Delta_j(v \cdot \nabla a)) \circ \psi_j, \\ \bar{\mathcal{R}}_j^2 &= \Lambda(d_j \circ \psi_j) - (\Lambda d_j) \circ \psi_j, \\ \bar{\mathcal{Q}}_j^1 &= ((S_{j-1} v \cdot \nabla d_j) - \Delta_j(v \cdot \nabla d)) \circ \psi_j, \\ \bar{\mathcal{Q}}_j^2 &= \bar{\nu}(\Delta d_j) \circ \psi_j - \bar{\nu} \Delta(d_j \circ \psi_j) + (\Lambda a_j) \circ \psi_j - \Lambda(a_j \circ \psi_j), \end{aligned}$$

Using Proposition 7.4 with $s = s_p, \sigma = s$ and Lemma 2.1, we get

$$\begin{aligned} &\|S_{j-1} v \cdot \nabla a_j - \Delta_j(v \cdot \nabla a)\|_{L_T^1 L^p} \\ &\leq \|[v, \Delta_j] \cdot \nabla a\|_{L_T^1 L^p} + \|(S_{j-1} v - v) \cdot \nabla a_j\|_{L_T^1 L^p} \\ &\leq C c(j) 2^{-sj} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{s_p, s}} + C \sum_{j' \geq j-1} 2^{j'} \|\Delta_{j'} v\|_{L_T^1 L^\infty} 2^{j-j'} \|a_j\|_{L_T^\infty L^p} \\ &\leq C c(j) 2^{-sj} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{s_p, s}}, \end{aligned}$$

from which, Lemma 2.2, and Lemma 2.7 (a), we deduce that

$$\begin{aligned} \|\Delta_k \bar{\mathcal{R}}_j^1\|_{L_T^1 L^p} &\leq C e^{C\bar{V}(T)} \|S_{j-1} v \cdot \nabla a_j - \Delta_j(v \cdot \nabla a)\|_{L_T^1 L^p} \\ &\leq C c(j) 2^{-sj} e^{C\bar{V}(T)} \bar{V}(T) \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{s_p-1,s}}. \end{aligned} \quad (5.10)$$

From Lemma 2.4, it follows that

$$\|\Delta_k \bar{\mathcal{R}}_j^2(t)\|_{L^p} \leq C 2^j e^{C\bar{V}(t)} \bar{V}(t)^{\frac{1}{2}} \|d_j\|_{L^p}. \quad (5.11)$$

Similarly, we have

$$\|\Delta_k \bar{\mathcal{Q}}_j^1\|_{L_T^1 L^p} \leq C c(j) 2^{-(s-1)j} e^{C\bar{V}(T)} \bar{V}(T) \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{s_p-1,s-1}}, \quad (5.12)$$

$$\|\Delta_k \bar{\mathcal{Q}}_j^2(t)\|_{L^p} \leq C e^{C\bar{V}(t)} \bar{V}(t)^{\frac{1}{2}} (2^{2j} \|d_j\|_{L^p} + 2^j \|a_j\|_{L^p}). \quad (5.13)$$

Now we are in position to estimate $\|(\Delta_k \bar{a}_j, \Delta_k \bar{d}_j)\|_{L_T^r L^p}$. Since $(\bar{a}_j, \bar{d}_j)$ satisfies (5.5), it can be expressed in terms of Green's matrix as

$$\begin{pmatrix} \Delta_k \bar{a}_j \\ \Delta_k \bar{d}_j \end{pmatrix} = \mathcal{G}(x, t) * \begin{pmatrix} \Delta_k a_j^0 \\ \Delta_k d_j^0 \end{pmatrix} + \int_0^t \mathcal{G}(x, t - \tau) * \begin{pmatrix} \Delta_k \bar{F}_j + \Delta_k \bar{\mathcal{R}}_j \\ \Delta_k \bar{G}_j + \Delta_k \bar{\mathcal{Q}}_j \end{pmatrix}(\tau) d\tau.$$

We consider two cases:

Case 1. $2^k > R_0$. By Lemma 4.2 (b), we have

$$\begin{aligned} \Delta_k \bar{a}_j(t, x) &= e^{-\bar{\nu}^{-1}t} \Delta_k a_j^0 + \mathcal{G}^1 * \Delta_k d_j^0 + \mathcal{G}^2 * \Delta_k a_j^0 \\ &\quad + \int_0^t \{e^{-\bar{\nu}^{-1}(t-\tau)} + \mathcal{G}^2(x, t - \tau) * \} (\Delta_k \bar{F}_j + \Delta_k \bar{\mathcal{R}}_j)(\tau) d\tau \\ &\quad + \int_0^t \mathcal{G}^1(x, t - \tau) * (\Delta_k \bar{G}_j + \Delta_k \bar{\mathcal{Q}}_j)(\tau) d\tau, \end{aligned}$$

where the scale function $\mathcal{G}^2$ corresponds to the first nonzero in the diagonal matrix (4.9). Noting that $2^k > R_0$, we apply Proposition 4.4 (c) to get

$$\begin{aligned} \|\Delta_k \bar{a}_j(t)\|_{L^p} &\leq C e^{-ct} \|\Delta_k a_j^0\|_{L^p} + C 2^{-k} e^{-ct} \|\Delta_k d_j^0\|_{L^p} \\ &\quad + C \int_0^t e^{-c(t-\tau)} (\|\Delta_k \bar{F}_j(\tau)\|_{L^p} + \|\Delta_k \bar{\mathcal{R}}_j(\tau)\|_{L^p}) d\tau \\ &\quad + C 2^{-k} \int_0^t e^{-c(t-\tau)} (\|\Delta_k \bar{G}_j(\tau)\|_{L^p} + \|\Delta_k \bar{\mathcal{Q}}_j(\tau)\|_{L^p}) d\tau. \end{aligned}$$

Taking L^r norm with respect to t and using Young's inequality give

$$\begin{aligned} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} &\leq C (\|\Delta_k a_j^0\|_{L^p} + 2^{-k} \|\Delta_k d_j^0\|_{L^p} + \|\Delta_k \bar{F}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{R}}_j\|_{L_T^1 L^p}) \\ &\quad + C 2^{-k} (\|\Delta_k \bar{G}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{Q}}_j\|_{L_T^1 L^p}). \end{aligned} \quad (5.14)$$

On the other hand, we have

$$\begin{aligned} \Delta_k \bar{d}_j(t, x) &= \mathcal{G}^1 * \Delta_k a_j^0 + e^{\bar{\nu} \Delta t} \Delta_k d_j^0 + \mathcal{G}^2 * \Delta_k d_j^0 \\ &\quad + \int_0^t \mathcal{G}^1(x, t - \tau) * (\Delta_k \bar{F}_j + \Delta_k \bar{\mathcal{R}}_j)(\tau) d\tau \\ &\quad + \int_0^t (e^{\bar{\nu} \Delta(t-\tau)} + \mathcal{G}^2(x, t - \tau) *) (\Delta_k \bar{G}_j + \Delta_k \bar{\mathcal{Q}}_j)(\tau) d\tau, \end{aligned}$$

where the scale function $\mathcal{G}^2$ corresponds to the second nonzero in the diagonal matrix (4.9). In analogy as leading to (5.14), we get

$$\begin{aligned} \|\Delta_k \bar{d}_j\|_{L_T^r L^p} &\leq C \left(2^{-k} \|\Delta_k a_j^0\|_{L^p} + 2^{-\frac{2}{r}k} \|\Delta_k d_j^0\|_{L^p} \right. \\ &\quad + 2^{-k} (\|\Delta_k \bar{F}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{R}}_j^1\|_{L_T^1 L^p}) + 2^{-k} \|\Delta_k \bar{\mathcal{R}}_j^2\|_{L_T^r L^p} \\ &\quad \left. + 2^{-\frac{2}{r}k} (\|\Delta_k \bar{G}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{Q}}_j^1\|_{L_T^1 L^p}) + 2^{-2k} \|\Delta_k \bar{\mathcal{Q}}_j^2\|_{L_T^r L^p} \right). \end{aligned} \quad (5.15)$$

Substituting (5.9)-(5.13) into (5.14) and (5.15), then summing over k , we obtain

$$\begin{aligned} \sum_{|k-j| \leq N, 2^k > R_0} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} &\leq C e^{C\bar{V}(T)} \left\{ \|a_j^0\|_{L^p} + 2^{-j} \|d_j^0\|_{L^p} \right. \\ &\quad + \bar{V}(T)^{\frac{1}{2}} (\|a_j\|_{L_T^1 L^p} + 2^j \|d_j\|_{L_T^1 L^p}) + \|F_j\|_{L_T^1 L^p} + 2^{-j} \|G_j\|_{L_T^1 L^p} \\ &\quad \left. + 2^{-sj} c(j) \bar{V}(T) (\|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s}} + \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1}}) \right\}, \end{aligned} \quad (5.16)$$

and

$$\begin{aligned} \sum_{|k-j| \leq N, 2^k > R_0} \|\Delta_k \bar{d}_j\|_{L_T^r L^p} &\leq C e^{C\bar{V}(T)} \left\{ 2^{-j} \|a_j^0\|_{L^p} + 2^{-\frac{2}{r}j} \|d_j^0\|_{L^p} \right. \\ &\quad + \bar{V}(T)^{\frac{1}{2}} (2^{-j} \|a_j\|_{L_T^r L^p} + \|d_j\|_{L_T^r L^p}) + 2^{-j} \|F_j\|_{L_T^1 L^p} + 2^{-\frac{2}{r}j} \|G_j\|_{L_T^1 L^p} \\ &\quad \left. + c(j) 2^{-(s-1+\frac{2}{r})j} \bar{V}(T) (\|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s}} + \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1}}) \right\}. \end{aligned} \quad (5.17)$$

Here we used $2^N \sim e^{C\bar{V}(T)}$ and the summation is finite (at most $2N+1$) for fixed j .

Case 2. $2^k \leq R_0$. Noting that

$$2^k \geq 2^{j-N} \geq R_0 2^{-N} \sim R_0 e^{-C\bar{V}(T)}.$$

Then we apply Proposition 4.4 (b) with $b \sim e^{-C\bar{V}(T)}$ to get

$$\begin{aligned} \|\Delta_k \bar{a}_j(t)\|_{L^p} + \|\Delta_k \bar{d}_j(t)\|_{L^p} &\leq C e^{C\bar{V}(T)} e^{-c2^{2k}t} (\|\Delta_k a_j^0\|_{L^p} + \|\Delta_k d_j^0\|_{L^p}) \\ &\quad + C e^{C\bar{V}(T)} \int_0^t e^{-c2^{2k}(t-\tau)} (\|\Delta_k \bar{F}_j(\tau)\|_{L^p} + \|\Delta_k \bar{G}_j(\tau)\|_{L^p} \\ &\quad + \|\Delta_k \bar{\mathcal{R}}_j(\tau)\|_{L^p} + \|\Delta_k \bar{\mathcal{Q}}_j(\tau)\|_{L^p}) d\tau. \end{aligned}$$

Then taking L^r norm with respect to t , we get by Young's inequality that

$$\begin{aligned} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} + \|\Delta_k \bar{d}_j\|_{L_T^r L^p} &\leq C e^{C\bar{V}(T)} \left\{ 2^{-\frac{2}{r}k} (\|\Delta_k a_j^0\|_{L^p} + \|\Delta_k d_j^0\|_{L^p} + \|\Delta_k \bar{F}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{R}}_j^1\|_{L_T^1 L^p} \right. \\ &\quad \left. + \|\Delta_k \bar{G}_j\|_{L_T^1 L^p} + \|\Delta_k \bar{\mathcal{Q}}_j^1\|_{L_T^1 L^p}) + 2^{-2k} (\|\Delta_k \bar{\mathcal{R}}_j^2\|_{L_T^r L^p} + \|\Delta_k \bar{\mathcal{Q}}_j^2\|_{L_T^r L^p}) \right\}. \end{aligned}$$

Substituting (5.9)-(5.13) into the above inequality, then summing over k , we obtain

$$\begin{aligned} \sum_{|k-j| \leq N, 2^k \leq R_0} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} + \|\Delta_k \bar{d}_j\|_{L_T^r L^p} &\leq C e^{C\bar{V}(T)} \left\{ 2^{-\frac{2}{r}j} (\|a_j^0\|_{L^p} + \|d_j^0\|_{L^p}) \right. \\ &\quad + 2^{-\frac{2}{r}j} (\|F_j\|_{L_T^1 L^p} + \|G_j\|_{L_T^1 L^p}) + \bar{V}(T)^{\frac{1}{2}} (2^{-j} \|a_j\|_{L_T^r L^p} + \|d_j\|_{L_T^r L^p}) \\ &\quad \left. + c(j) \bar{V}(T) (2^{-(s+\frac{2}{r})j} \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s}} + 2^{-(s-1+\frac{2}{r})j} \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1}}) \right\}. \end{aligned} \quad (5.18)$$

Summing up (5.16)–(5.18) and noting that for $2^k \leq R_0$,

$$R_0 < 2^j \leq 2^{k+N} \leq R_0 2^N \sim R_0 e^{C\bar{V}(T)},$$

we deduce that for any $1 \leq r \leq \infty$,

$$\begin{aligned} & \sum_{|k-j| \leq N} 2^{js} \|\Delta_k \bar{a}_j\|_{L_T^r L^p} + 2^{j(s-1+\frac{2}{r})} \|\Delta_k \bar{d}_j\|_{L_T^r L^p} \\ & \leq C e^{C\bar{V}(T)} \left\{ 2^{js} (\|a_j^0\|_{L^p} + 2^{-j} \|d_j^0\|_{L^p}) \right. \\ & \quad + \bar{V}(T)^{\frac{1}{2}} (2^{j(s-1+\frac{2}{r})} \|d_j\|_{L_T^r L^p} + 2^{js} \|a_j\|_{L_T^r L^p \cap L_T^1 L^p} + 2^{j(s+1)} \|d_j\|_{L_T^1 L^p}) \\ & \quad \left. + 2^{js} \|F_j\|_{L_T^1 L^p} + 2^{j(s-1)} \|G_j\|_{L_T^1 L^p} + c(j) \bar{V}(T) \|(a, d)\|_{\mathcal{E}_T^s} \right\}, \end{aligned}$$

which together with (5.7) and (5.8) implies that

$$\begin{aligned} & \sum_{2^j > R_0} (2^{js} \|a_j\|_{L_T^\infty L^p} + 2^{js} \|a_j\|_{L_T^1 L^p} + 2^{j(s-1)} \|d_j\|_{L_T^\infty L^p} + 2^{j(s+1)} \|d_j\|_{L_T^1 L^p}) \\ & \leq C e^{C\bar{V}(T)} \left\{ \sum_{2^j > R_0} 2^{js} (\|a_j^0\|_{L^p} + 2^{-j} \|d_j^0\|_{L^p}) + \|F\|_{L_T^1 \dot{B}_{2,p}^{sp-1,s}} + \|G\|_{L_T^1 \dot{B}_{2,p}^{sp-1,s-1}} \right. \\ & \quad \left. + (\bar{V}(T) + \bar{V}(T)^{\frac{1}{2}}) \|(a, d)\|_{\mathcal{E}_T^s} \right\}. \end{aligned} \quad (5.19)$$

5.3. Proof of Theorem 5.1. Thanks to (5.3) and (5.19), we obtain

$$\begin{aligned} & \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s} \cap L_T^1 \dot{B}_{2,p}^{sp+1,s}} + \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1} \cap L_T^1 \dot{B}_{2,p}^{sp+1,s+1}} \\ & \leq C e^{C\bar{V}(T)} \left\{ \|a_0\|_{\dot{B}_{2,p}^{sp-1,s}} + \|d_0\|_{\dot{B}_{2,p}^{sp-1,s-1}} + \|F\|_{L_T^1 \dot{B}_{2,p}^{sp-1,s}} + \|G\|_{L_T^1 \dot{B}_{2,p}^{sp-1,s-1}} \right. \\ & \quad + \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \left(\|a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p} + \frac{n}{p'}, s}} + \|d\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p} + \frac{n}{p'}, sp}} \right) \\ & \quad \left. + \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \left(\|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{sp,s}} + \|d\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{sp,s}} \right) + (\bar{V}(T) + \bar{V}(T)^{\frac{1}{2}}) \|(a, d)\|_{\mathcal{E}_T^s} \right\}. \end{aligned}$$

This together with the following interpolations inequalities

$$\begin{aligned} & \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \leq \|v\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}^{\frac{1}{2}} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}^{\frac{1}{2}}, \\ & \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{p}, \frac{2n}{p} - \frac{n}{2}}} \leq \|v\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}^{\frac{1}{\tilde{p}'}} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}^{\frac{1}{\tilde{p}}}, \\ & \|d\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p} + \frac{n}{p'}, sp}} \leq \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1}}^{\frac{1}{\tilde{p}}} \|d\|_{L_T^1 \dot{B}_{2,p}^{sp+1,s+1}}^{\frac{1}{\tilde{p}'}} , \\ & \|d\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{sp,s}} \leq \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s-1}}^{\frac{1}{2}} \|d\|_{L_T^1 \dot{B}_{2,p}^{sp+1,s+1}}^{\frac{1}{2}}, \\ & \|a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{s-\frac{n}{p} + \frac{n}{p'}, s}} \leq \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s}}^{\frac{1}{\tilde{p}}} \|a\|_{L_T^1 \dot{B}_{2,p}^{sp+1,s}}^{\frac{1}{\tilde{p}'}} , \\ & \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{sp,s}} \leq \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{sp-1,s}}^{\frac{1}{2}} \|a\|_{L_T^1 \dot{B}_{2,p}^{sp+1,s}}^{\frac{1}{2}}, \end{aligned}$$

and Young's inequality yield Theorem 5.1. $\square$

5.4. Proof of Theorem 5.2. Since the proof is essentially the same as that of Theorem 5.1, here we only present main differences. We use Proposition 7.3 (c) to obtain

$$\begin{aligned} \sum_{j \in \mathbb{Z}} 2^{j(s-1)} \|\Delta_j(v \nabla a)\|_{L_T^1 L^2} &\leq C(\|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s} + \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s}), \\ \sum_{j \in \mathbb{Z}} 2^{j(s-1)} \|\Delta_j(v \nabla d)\|_{L_T^1 L^2} &\leq C(\|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|d\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s} + \|d\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s}). \end{aligned}$$

Then in the case of low frequency, we can conclude that

$$\begin{aligned} &\sum_{2^j \leq R_0} 2^{j(s-1+\frac{2}{r})} (\|a_j\|_{L_T^r L^2} + \|d_j\|_{L_T^r L^2}) \\ &\leq C \left(\sum_{2^j \leq R_0} 2^{j(s-1)} (\|a_j^0\|_{L^2} + \|d_j^0\|_{L^2}) + \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|(a, d)\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s} \right. \\ &\quad \left. + \|v\|_{\tilde{L}_T^2 \dot{B}_{2,1}^s} \|(a, d)\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} + \|F\|_{L_T^1 \dot{B}^{s-1, s}} + \|G\|_{L_T^1 \dot{B}_{2,1}^{s-1}} \right). \end{aligned} \quad (5.20)$$

From Proposition 7.4 and Lemma 2.2, we infer that

$$\begin{aligned} \|\Delta_k \bar{\mathcal{R}}_j^1\|_{L_T^1 L^2} &\leq Cc(j) 2^{-sj} e^{\bar{V}(T)} \bar{V}(T) \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,1}^s}, \\ \|\Delta_k \bar{\mathcal{Q}}_j^1\|_{L_T^1 L^2} &\leq Cc(j) 2^{-(s-1)j} e^{\bar{V}(T)} \bar{V}(T) \|d\|_{\tilde{L}_T^\infty \dot{B}_{2,1}^{s-1}}. \end{aligned}$$

Then in the case of high frequency, we can conclude that

$$\begin{aligned} &\sum_{2^j > R_0} (2^{js} \|a_j\|_{L_T^\infty L^2} + 2^{js} \|a_j\|_{L_T^1 L^2} + 2^{j(s-1)} \|d_j\|_{L_T^\infty L^2} + 2^{j(s+1)} \|d_j\|_{L_T^1 L^2}) \\ &\leq C e^{C\bar{V}(T)} \left\{ \sum_{2^j > R_0} 2^{js} (\|a_j^0\|_{L^2} + 2^{-j} \|d_j^0\|_{L^2}) + \|F\|_{L_T^1 \dot{B}^{s-1, s}} + \|G\|_{L_T^1 \dot{B}_{2,1}^{s-1}} \right. \\ &\quad \left. + (\bar{V}(T) + \bar{V}(T)^{\frac{1}{2}}) \|(a, d)\|_{E_T^s} \right\}. \end{aligned}$$

This together with (5.20) implies Theorem 5.2. $\square$

6. PROOF OF THEOREM 1.1

This section is devoted to the proof of Theorem 1.1. We denote

$$a(t, x) \stackrel{\text{def}}{=} \frac{\rho(\varpi^{-2}t, \varpi^{-1}x)}{\bar{\rho}} - 1, \quad v(t, x) \stackrel{\text{def}}{=} \varpi^{-1}u(\varpi^{-2}t, \varpi^{-1}x),$$

where $\varpi \stackrel{\text{def}}{=} (P'(\bar{\rho}))^{\frac{1}{2}}$. Then the system (1.1) can be rewritten as

$$\begin{cases} \partial_t a + v \cdot \nabla a + \operatorname{div} v = -a \operatorname{div} v, \\ \partial_t v + v \cdot \nabla v - \mathcal{A}v + \nabla a = -L(a)\mathcal{A}v - K(a)\nabla a, \\ (a, v)|_{t=0} = (a_0, v_0), \end{cases} \quad (6.1)$$

where

$$\mathcal{A} = \bar{\mu} \Delta + (\bar{\lambda} + \bar{\mu}) \nabla \operatorname{div}, \quad K(a) = \frac{P'(\bar{\rho}(1+a))}{(1+a)P'(\bar{\rho})} - 1 \quad \text{and} \quad L(a) = \frac{a}{1+a},$$

with $\bar{\mu} = \frac{\mu}{\rho}$ and $\bar{\lambda} = \frac{\lambda}{\rho}$. We denote

$$d \stackrel{\text{def}}{=} \Lambda^{-1} \operatorname{div} v, \quad \Omega \stackrel{\text{def}}{=} \Lambda^{-1} (\operatorname{curl} v)_j^i.$$

In view of the equality

$$\Delta v^i = \partial_i \operatorname{div} v + \partial_j (\operatorname{curl} v)_j^i, \quad \text{where } (\operatorname{curl} v)_j^i = \partial_j v^i - \partial_i v^j,$$

we find that (a, d, Ω) satisfies

$$\begin{cases} \partial_t a + \Lambda d + v \cdot \nabla a = F, \\ \partial_t d - \bar{\nu} \Delta d - \Lambda a + v \cdot \nabla d = G, \\ \partial_t \Omega - \bar{\mu} \Delta \Omega = H, \\ (a, d, \Omega)|_{t=0} = (a_0, \Lambda^{-1} \operatorname{div} v_0, \Lambda^{-1} \operatorname{curl} v_0), \end{cases} \quad (6.2)$$

where $\bar{\nu} = 2\bar{\mu} + \bar{\lambda}$ and

$$\begin{aligned} F &= -a \operatorname{div} v, \\ G &= v \cdot \nabla d - \Lambda^{-1} \operatorname{div}(v \cdot \nabla v + L(a) \mathcal{A}v + K(a) \nabla a), \\ H &= -\Lambda^{-1} \operatorname{curl}(v \cdot \nabla v + L(a) \mathcal{A}v), \\ v &= -\Lambda^{-1} \nabla d - \Lambda^{-1} \operatorname{div} \Omega. \end{aligned}$$

6.1. A priori estimates.

Proposition 6.1. Let $2 \leq p < 2n$, $p \leq \min(4, \frac{2n}{n-2})$. Assume that (a, d, Ω) is a smooth solution of the system (6.2) on $[0, T]$ with

$$\|a\|_{L^\infty([0, T] \times \mathbb{R}^n)} \leq \frac{1}{2}.$$

Then we have

$$\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} \leq C e^{C\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}} \left\{ \|(a_0, v_0)\|_{\mathcal{E}_0^{\frac{n}{p}}} + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}^{\frac{3}{2}} (1 + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}})^{\frac{n}{2}+2} \right\}. \quad (6.3)$$

Here $\|(a_0, v_0)\|_{\mathcal{E}_0^{\frac{n}{p}}} \stackrel{\text{def}}{=} \|a_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} + \|v_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}$.

Proof. Thanks to Theorem 5.1, we get

$$\begin{aligned} \|(a, d)\|_{\mathcal{E}_T^{\frac{n}{p}}} &\leq C e^{C\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}} \left\{ \|(a_0, d_0)\|_{\mathcal{E}_0^{\frac{n}{p}}} + (\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}^{\frac{1}{2}} + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}) \right. \\ &\quad \left. \times \|(a, d)\|_{\mathcal{E}_T^{\frac{n}{p}}} + \|F\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} + \|G\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \right\}, \end{aligned}$$

and from Proposition 3.2,

$$\|\Omega\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} + \|\Omega\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \leq C(\|\Omega_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} + \|H\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}}).$$

Then we obtain

$$\begin{aligned} \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} &\leq C e^{C\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}} \left\{ \|(a_0, v_0)\|_{\mathcal{E}_0^{\frac{n}{p}}} + (\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}^{\frac{1}{2}} + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}) \right. \\ &\quad \left. \times \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} + \|F\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} + \|(G, H)\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \right\}. \quad (6.4) \end{aligned}$$

Here we used the fact

$$\|v\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s, \sigma}} \approx \|d\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s, \sigma}} + \|\Omega\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s, \sigma}}$$

for any $s, \sigma \in \mathbb{R}, r \in [1, \infty]$.

Now we need to estimate the nonlinear terms F, G and H in (6.4). We define $\tilde{p}$ as $\frac{1}{\tilde{p}} = \frac{n}{2p} - \frac{n}{4} + \frac{1}{2}$, and denote $\tilde{p}'$ by the conjugate index of $\tilde{p}$. Due to $2 \leq p < 2n$, $p \leq \frac{2n}{n-2}$, using Proposition 7.3 (b) with $s = \frac{n}{p}, t = \frac{n}{p'} - 1, \tilde{s} = \frac{n}{2} - 1, \tilde{t} = \frac{n}{2}, \gamma = 0$ yields

$$\begin{aligned} & \sum_{2^j \leq R_0} 2^{j(\frac{n}{2}-1)} \|\Delta_j F\|_{L_T^1 L^2} \\ & \leq C \|a\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{2n}{p} - \frac{n}{2}}} \|\operatorname{div} v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1}} + C \|\operatorname{div} v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \\ & \leq C \|a\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{2}}} + C \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}}, \end{aligned}$$

and from Proposition 7.3 (a) with $\sigma = \tau = \frac{n}{p}$,

$$\sum_{2^j > R_0} 2^{j\frac{n}{p}} \|\Delta_j F\|_{L_T^1 L^p} \leq C \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|\operatorname{div} v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \leq C \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}},$$

from which, we infer that

$$\begin{aligned} \|F\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} & \leq C \left(\|a\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{2}}} + \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \right. \\ & \quad \left. + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \right). \end{aligned} \quad (6.5)$$

Similarly, we have

$$\begin{aligned} & \|v \nabla v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} + \|v \nabla d\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \\ & \leq C \|v\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{2n}{p} - \frac{n}{2}}} \|v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{2}}} + C \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}}. \end{aligned}$$

From Proposition 7.3 (b) with $\gamma = \frac{n}{p} - \frac{n}{2}$ and Proposition 7.5, it follows that

$$\begin{aligned} & \sum_{2^j \leq R_0} 2^{j(\frac{n}{2}-1)} \|\Delta_j (K(a) \nabla a)\|_{L_T^1 L^2} \\ & \leq C \|K(a)\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{2n}{p} - \frac{n}{2}}} \|\nabla a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1}} + C \|\nabla a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1}} \|K(a)\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \\ & \leq C \left(\|a\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{2}}} + \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \right) (1 + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}})^{\frac{n}{2}+1}, \end{aligned}$$

and with $\gamma = -1$

$$\begin{aligned} & \sum_{2^j \leq R_0} 2^{j(\frac{n}{2}-1)} \|\Delta_j (L(a) \mathcal{A}v)\|_{L_T^1 L^2} \\ & \leq C \|L(a)\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{2n}{p} - \frac{n}{2}}} \|\mathcal{A}v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1+\gamma}} + C \|\mathcal{A}v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{2}-1}} \|L(a)\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \\ & \leq C \left(\|a\|_{\tilde{L}_T^{\tilde{p}} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{\tilde{L}_T^{\tilde{p}'} \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{2}}} + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \right) (1 + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}})^{\frac{n}{2}+1}. \end{aligned}$$

On the other hand, from Proposition 7.3 (a) and Proposition 7.5,

$$\begin{aligned}
& \sum_{2^j > R_0} 2^{j(\frac{n}{p}-1)} (\|\Delta_j(K(a)\nabla a)\|_{L_T^1 L^p} + \|\Delta_j(L(a)\mathcal{A}v)\|_{L_T^1 L^p}) \\
& \leq C\|K(a)\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|\nabla a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} + C\|L(a)\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|Av\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \\
& \leq C(1 + \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\frac{n}{2}+1} (\|a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} + \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}).
\end{aligned}$$

Thus, we deduce that

$$\begin{aligned}
\|(G, H)\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} & \leq C(1 + \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\frac{n}{2}+1} \left(\|a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \right. \\
& \quad \left. + \|a\|_{\tilde{L}_T^p \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} (\|a\|_{\tilde{L}_T^{\frac{p'}{p}} \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} + \|v\|_{\tilde{L}_T^{\frac{p'}{p}} \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}}) + \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|v\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \right). \quad (6.6)
\end{aligned}$$

Noting that the interpolation inequalities

$$\begin{aligned}
\|a\|_{\tilde{L}_T^p \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} & \leq \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}}^{\frac{1}{p'}} \|a\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}}}^{\frac{1}{p}}, \\
\|a\|_{\tilde{L}_T^{\frac{p'}{p}} \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} & \leq \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}}^{\frac{1}{p'}} \|a\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}}}^{\frac{1}{p}}, \\
\|a\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} & \leq \|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}}^{\frac{1}{2}} \|a\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}}}^{\frac{1}{2}}, \\
\|v\|_{\tilde{L}_T^{\frac{p'}{p}} \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} & \leq \|v\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}^{\frac{1}{p'}} \|v\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}^{\frac{1}{p}}, \\
\|v\|_{\tilde{L}_T^p \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{2n}{p}-\frac{n}{2}}} & \leq \|v\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}^{\frac{1}{p'}} \|v\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}^{\frac{1}{p}}, \\
\|v\|_{\tilde{L}_T^2 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} & \leq \|v\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}}^{\frac{1}{2}} \|v\|_{L_T^1 \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}^{\frac{1}{2}},
\end{aligned}$$

and from $p \leq \frac{2n}{n-2}$,

$$\|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} \leq C\|a\|_{\tilde{L}_T^\infty \dot{\mathcal{B}}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}},$$

we plug (6.5)-(6.6) into (6.4) to obtain (6.3). $\square$

Proposition 6.2. Under the assumption of Proposition 6.1, we have

$$\begin{aligned}
\|(a, v)\|_{E_T^{\frac{n}{2}}} & \leq C e^{C\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}} \left\{ \|(a_0, v_0)\|_{E_0^{\frac{n}{2}}} + \|(a, v)\|_{E_T^{\frac{n}{2}}} \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}^{\frac{1}{2}} \right. \\
& \quad \left. \times (1 + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}})^{\frac{n}{2}+2} \right\}. \quad (6.7)
\end{aligned}$$

Here $\|(a_0, v_0)\|_{E_0^{\frac{n}{2}}} \stackrel{\text{def}}{=} \|a_0\|_{\dot{\mathcal{B}}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}} + \|v_0\|_{\dot{\mathcal{B}}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}}$.

Proof. Since the proof is similar to that of Proposition 6.1, here we only indicate main differences. Firstly, we get by using Theorem 5.2 and Proposition 3.2 that

$$\begin{aligned} \|(a, v)\|_{E_T^{\frac{n}{2}}} &\leq C e^{C\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}} \left\{ \|(a_0, v_0)\|_{E_0^{\frac{n}{2}}} + \left(\|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}}^{\frac{1}{2}} + \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} \right) \|(a, v)\|_{E_T^{\frac{n}{2}}} \right. \\ &\quad \left. + \|F\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}} + \|(G, H)\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1}} \right\}. \end{aligned} \quad (6.8)$$

Applying Proposition 7.3 (c) yields

$$\begin{aligned} \sum_{2^j \leq R_0} 2^{j(\frac{n}{2}-1)} \|\Delta_j F\|_{L_T^1 L^2} &\leq C \|a\|_{L_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \|\operatorname{div} v\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}}} + C \|\operatorname{div} v\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,1}^{\frac{n}{2}}}, \\ \sum_{2^j > R_0} 2^{j\frac{n}{2}} \|\Delta_j F\|_{L_T^1 L^2} &\leq C \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|\operatorname{div} v\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}}} + C \|\operatorname{div} v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,1}^{\frac{n}{2}}}, \end{aligned}$$

which imply by the interpolation and Lemma 2.7 (a) that

$$\|F\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1, \frac{n}{2}}} \leq C \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} \|(a, v)\|_{E_T^{\frac{n}{2}}}. \quad (6.9)$$

Similarly, we have

$$\begin{aligned} &\sum_{j \in \mathbb{Z}} 2^{j(\frac{n}{2}-1)} (\|\Delta_j(v \nabla d)\|_{L_T^1 L^2} + \|\Delta_j(v \nabla v)\|_{L_T^1 L^2}) \\ &\leq C \|v\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \|v\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}+1}} + C \|v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|v\|_{\tilde{L}_T^\infty \dot{B}_{2,1}^{\frac{n}{2}-1}}. \end{aligned}$$

From Proposition 7.3 (c) and Proposition 7.5, we infer that

$$\begin{aligned} &\sum_{j \in \mathbb{Z}} 2^{j(\frac{n}{2}-1)} \|\Delta_j(K(a) \nabla a)\|_{L_T^1 L^2} \\ &\leq C \|K(a)\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|\nabla a\|_{\tilde{L}_T^2 \dot{B}_{2,1}^{\frac{n}{2}-1}} + C \|\nabla a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \|K(a)\|_{\tilde{L}_T^2 \dot{B}_{2,1}^{\frac{n}{2}}} \\ &\leq C (1 + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\frac{n}{2}+1} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|a\|_{\tilde{L}_T^2 \dot{B}_{2,1}^{\frac{n}{2}}}, \end{aligned}$$

and

$$\begin{aligned} &\sum_{j \in \mathbb{Z}} 2^{j(\frac{n}{2}-1)} \|\Delta_j(L(a) \mathcal{A}v)\|_{L_T^1 L^2} \\ &\leq C \|L(a)\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}}} \|\mathcal{A}v\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1}} + C \|\mathcal{A}v\|_{L_T^1 \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \|L(a)\|_{\tilde{L}_T^\infty \dot{B}_{2,1}^{\frac{n}{2}}} \\ &\leq C (1 + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\frac{n}{2}+1} \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} \|(a, v)\|_{E_T^{\frac{n}{2}}}. \end{aligned}$$

Thus, we obtain

$$\|G\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1}} + \|H\|_{L_T^1 \dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C (1 + \|a\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\frac{n}{2}+1} \|(a, v)\|_{\mathcal{E}_T^{\frac{n}{p}}} \|(a, v)\|_{E_T^{\frac{n}{2}}}.$$

This together with (6.8) and (6.9) implies (6.7). $\square$

6.2. Approximate solutions and uniform estimates. The construction of approximate solutions is based on the following local existence theorem.

Theorem 6.3. [15] Assume that $\rho_0 - \bar{\rho} \in \dot{B}_{2,1}^{\frac{n}{2}}$ and $u_0 \in \dot{B}_{2,1}^{\frac{n}{2}-1}$ with ρ_0 bounded away from zero. There exists a positive time T such that the system (1.1) has a unique solution (ρ, u) with ρ bounded away from zero,

$$\rho - \bar{\rho} \in C([0, T]; \dot{B}_{2,1}^{\frac{n}{2}}) \quad \text{and} \quad u \in C([0, T]; \dot{B}_{2,1}^{\frac{n}{2}-1}) \cap L^1(0, T; \dot{B}_{2,1}^{\frac{n}{2}+1}).$$

Moreover, the solution (ρ, u) can be continued beyond T if the following three conditions hold:

- (a) The function $\rho - \bar{\rho}$ belongs to $L^\infty(0, T; \dot{B}_{2,1}^{\frac{n}{2}})$;
- (b) The function ρ is bounded away from zero;
- (c) $\int_0^T \|\nabla u(\tau)\|_{L^\infty} d\tau < \infty$.

To apply Theorem 6.3, we need

Lemma 6.4. Let $p \geq 2$. For any $(\rho_0 - \bar{\rho}, u_0) \in \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}} \times (\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1})^n$ satisfying $\rho_0 \geq \bar{\rho}$, there exists a sequence $\{(\rho_0^k, u_0^k)\}_{k \in \mathbb{N}}$ with $(\rho_0^k - \bar{\rho}, u_0^k) \in \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}} \times (\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1})^n$ such that

$$\|\rho_0^k - \rho_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}} \longrightarrow 0, \quad \|u_0^k - u_0\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}} \longrightarrow 0 \quad (6.10)$$

as $k \rightarrow \infty$, and $\rho_{0,k} \geq \frac{\bar{\rho}}{2}$ for any $k \in \mathbb{N}$.

Proof. This lemma can be proved by following the proof of Lemma 4.2 in [1]. $\square$

Let (ρ_0^k, u_0^k) be as stated in Lemma 6.4. Then Theorem 6.3 ensures that there exists a maximal existence time $T_k > 0$ such that the system (1.1) with the initial data $(\rho_{0,k}, u_{0,k})$ has a unique solution (ρ^k, u^k) with ρ^k bounded away from zero,

$$\rho^k - \bar{\rho} \in C([0, T_k]; \dot{B}_{2,1}^{\frac{n}{2}}) \quad \text{and} \quad u^k \in C([0, T_k]; \dot{B}_{2,1}^{\frac{n}{2}-1}) \cap L^1(0, T_k; \dot{B}_{2,1}^{\frac{n}{2}+1}).$$

Using the definition of the Besov space and Lemma 2.1, it is easy to check that

$$\rho^k - \bar{\rho} \in C([0, T_k]; \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}) \quad \text{and} \quad u^k \in C([0, T_k]; \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1}) \cap L^1(0, T_k; \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}).$$

We set

$$a^k(t, x) = \frac{\rho^k(\varpi^{-2}t, \varpi^{-1}x)}{\bar{\rho}} - 1, \quad v^k(t, x) = \varpi^{-1}u^k(\varpi^{-2}t, \varpi^{-1}x).$$

Then from (1.10) and (6.10), we find

$$\|(a_0^k, v_0^k)\|_{\mathcal{E}_0^{\frac{n}{p}}} \leq C_0 \eta,$$

for some constant C_0 . Given a constant M to be chosen later on, let us define

$$T_k^* \stackrel{\text{def}}{=} \sup \left\{ t \in [0, T_k]; \|(a^k, v^k)\|_{\mathcal{E}_t^{\frac{n}{p}}} \leq M\eta \right\}.$$

Firstly, we claim that

$$T_k^* = T_k, \quad \forall k \in \mathbb{N}.$$

Using the continuity argument, it suffices to show that for all $k \in \mathbb{N}$,

$$\|(a^k, v^k)\|_{\mathcal{E}_{T_k^*}^{\frac{n}{p}}} \leq \frac{3}{4} M\eta. \quad (6.11)$$

In fact, noticing that $\|a^k\|_{L^\infty} \leq C_1 \|a^k\|_{\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}}$, we choose η as

$$M\eta \leq \frac{1}{2C_1}$$

such that

$$\|a^k\|_{L^\infty([0, T_k^*) \times \mathbb{R}^n)} \leq \frac{1}{2}.$$

Thus, we can apply Proposition 6.1 to obtain

$$\|(a^k, v^k)\|_{\mathcal{E}_{T_k^*}^{\frac{n}{p}}} \leq Ce^{CM\eta} \left\{ C_0\eta + (M\eta)^{\frac{3}{2}}(1 + M\eta)^{\frac{n}{2}+2} \right\}. \quad (6.12)$$

Set $M = 4CC_0$, then choose η small enough such that

$$e^{CM\eta} \leq \frac{3}{2}, \quad C(M\eta)^{\frac{1}{2}}(1 + M\eta)^{\frac{n}{2}+2} \leq \frac{1}{4},$$

then the inequality (6.11) follows from (6.12). In conclusion, we construct a sequence of approximate solution (ρ^k, u^k) of (1.1) on $[0, T_k)$ satisfying

$$\|(a^k, v^k)\|_{\mathcal{E}_{T_k}^{\frac{n}{p}}} \leq M\eta, \quad (6.13)$$

for any $k \in \mathbb{N}$. Next, we claim that

$$T_k = +\infty, \quad \forall k \in \mathbb{N}.$$

Thanks to Theorem 6.3 and (6.13), it remains to prove $a^k \in L^\infty(0, T_k; \dot{B}_{2,1}^{\frac{n}{2}})$. While thanks to Proposition 6.2 and (6.13), we have

$$\|(a^k, v^k)\|_{E_{T_k}^{\frac{n}{2}}} \leq Ce^{CM\eta} \left\{ \|(a_0^k, v_0^k)\|_{E_0^{\frac{n}{2}}} + \|(a^k, v^k)\|_{E_{T_k}^{\frac{n}{2}}} (M\eta)^{\frac{1}{2}}(1 + M\eta)^{\frac{n}{2}+2} \right\},$$

which implies that

$$\|(a^k, v^k)\|_{E_{T_k}^{\frac{n}{2}}} \leq C \|(a_0^k, v_0^k)\|_{E_0^{\frac{n}{2}}}.$$

6.3. Existence. We will use the compact argument to prove the existence of the solution. Due to (6.13), it is easy to verify that

- a^k is uniformly bounded in $L^\infty(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}})$;
- v^k is uniformly bounded in $L^\infty(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}-1}) \cap L^1(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}+1})$.

By the interpolation, we also have

$$v^k \text{ is uniformly bounded in } L^{\frac{2}{1-\epsilon}}(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}-\epsilon}),$$

for any $\epsilon \in [-1, 1]$.

Let v_L^k be a solution of

$$\partial_t v_L^k - \mathcal{A}v_L^k = 0, \quad v_L^k(0) = v_0^k.$$

It is easy to check that v_L^k tends to the solution of

$$\partial_t v_L - \mathcal{A}v_L = 0, \quad v_L(0) = v_0,$$

in $L^\infty(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}-1}) \cap L^1(0, \infty; \dot{B}_{p,1}^{\frac{n}{p}+1})$.

We set $\tilde{a}^k \stackrel{\text{def}}{=} a^k - a_0^k$ and $\tilde{v}^k \stackrel{\text{def}}{=} v^k - v_L^k$. Firstly, we claim that $(\tilde{a}^k, \tilde{v}^k)$ is uniformly bounded in $C_{loc}^{\frac{1}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1}) \times C_{loc}^{\frac{2-\varsigma}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma})$ with

$$\varsigma = \min\left(\frac{2n}{p} - 1, 1\right).$$

Recall that

$$\partial_t \tilde{a}^k = -v^k \cdot \nabla a^k - \operatorname{div} v^k - a^k \operatorname{div} v^k,$$

which together with Lemma 2.8 implies $\partial_t \tilde{a}^k \in L^2 \dot{B}_{p,1}^{\frac{n}{p}-1}$, thus $\tilde{a}^k$ is uniformly bounded in $C^{\frac{1}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1})$. On the other hand,

$$\partial_t \tilde{v}^k = -v^k \cdot \nabla v^k + \mathcal{A} \tilde{v}^k - \nabla a^k - L(a^k) \mathcal{A} v^k - K(a^k) \nabla a^k.$$

Thanks to Lemma 2.8 and Proposition 7.5, we have

$$\begin{aligned} & \|v^k \cdot \nabla v^k + L(a^k) \mathcal{A} v^k\|_{L^{\frac{2}{2-\varsigma}} \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}} \\ & \leq C \|v^k\|_{L^2 \dot{B}_{p,1}^{\frac{n}{p}}} \|\nabla v^k\|_{L^{\frac{2}{1-\varsigma}} \dot{B}_{p,1}^{\frac{n}{p}-\varsigma-1}} + C(\|a^k\|_{L^\infty \dot{B}_{p,1}^{\frac{n}{p}}}) \|\mathcal{A} v^k\|_{L^{\frac{2}{2-\varsigma}} \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}}. \end{aligned}$$

The inclusion map $\dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}} \hookrightarrow \dot{B}_{p,1}^{\frac{n}{p}-\varsigma}$ implies $\Lambda a^k \in L^\infty \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}$, thus $K(a^k) \nabla a^k$ is bounded in $L^\infty \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}$. Therefore, we have $\partial_t \tilde{v}^k \in L_{loc}^{\frac{2}{2-\varsigma}} \dot{B}_p^{\frac{n}{p}-1-\varsigma}$, thus $\tilde{v}^k$ is uniformly bounded in $C_{loc}^{\frac{2-\varsigma}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma})$.

Let $\{\chi_j\}_{j \in \mathbb{N}}$ be a sequence of smooth functions supported in the ball $B(0, j+1)$ and equal to 1 on $B(0, j)$. The claim ensures that for any $j \in \mathbb{N}$, $\{\chi_j \tilde{a}^k\}_{k \in \mathbb{N}}$ is uniformly bounded in $C_{loc}^{\frac{1}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1})$, and $\{\chi_j \tilde{v}^k\}_{k \in \mathbb{N}}$ is uniformly bounded in $C_{loc}^{\frac{2-\varsigma}{2}}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma})$. Observe that for any $\chi \in C_0^\infty(\mathbb{R}^n)$, the map: $(\tilde{a}^k, \tilde{v}^k) \mapsto (\chi \tilde{a}^k, \chi \tilde{v}^k)$ is compact from

$$(\dot{B}_{p,1}^{\frac{n}{p}-1} \cap \dot{B}_{p,1}^{\frac{n}{p}}) \times (\dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma} \cap \dot{B}_{p,1}^{\frac{n}{p}-1}) \quad \text{into} \quad \dot{B}_{p,1}^{\frac{n}{p}-1} \times \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}.$$

By applying Ascoli's theorem and Cantor's diagonal process, there exists some function $(\tilde{a}, \tilde{v}) \in \mathcal{E}^{\frac{n}{p}}$ such that for any $j \in \mathbb{N}$,

$$\begin{aligned} \chi_j \tilde{a}^k & \longrightarrow \chi_j \tilde{a} \quad \text{in} \quad C_{loc}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1}), \\ \chi_j \tilde{v}^k & \longrightarrow \chi_j \tilde{v} \quad \text{in} \quad C_{loc}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{n}{p}-1-\varsigma}), \end{aligned} \tag{6.14}$$

as k tends to ∞ (up to a subsequence). By the interpolation, we also have

$$\begin{aligned} \chi_j a^k & \longrightarrow \chi_j \tilde{a} \quad \text{in} \quad C_{loc}(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{N}{p}-s}), \quad \forall 0 < s \leq 1, \\ \chi_j v^k & \longrightarrow \chi_j \tilde{v} \quad \text{in} \quad L_{loc}^1(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{N}{p}+s}), \quad \forall -1 \leq s < 1. \end{aligned} \tag{6.15}$$

With (6.14)-(6.15), it is a routine process to verify that $(\tilde{a} + a_0, \tilde{v} + v_L)$ satisfies the system (6.1) in the sense of distribution (see also [11]). Finally, following the argument in [11], we can show that $(a, v) \in C([0, \infty); \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}}) \times C([0, \infty); \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{n}{p}-1})$.

6.4. Uniqueness. In this subsection, we prove the uniqueness of the solution. Assume that $(\rho^1, u^1) \in \mathcal{E}^{\frac{n}{p}}$ and $(\rho^2, u^2) \in \mathcal{E}^{\frac{n}{p}}$ are two solutions of the system (1.1) with the same initial data. Without loss of generality, we may assume that (ρ^1, u^1) satisfies

$$\|(\rho^1 - \bar{\rho}, u^1)\|_{\mathcal{E}^{\frac{n}{p}}} \leq M\eta. \quad (6.16)$$

Since $\rho^2 - \bar{\rho} \in C([0, T]; \dot{B}_{2,p}^{\frac{n}{2}-1, \frac{N}{p}})$ and $\rho^2(0, x) \geq \frac{\bar{\rho}}{2}$, there exists a positive time T such that

$$\rho^2(t, x) \geq \frac{\bar{\rho}}{3}, \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n.$$

We set

$$\begin{aligned} a^k(t, x) &= \frac{\rho^k(\varpi^{-2}t, \varpi^{-1}x)}{\bar{\rho}} - 1, \quad v^k(t, x) = \varpi^{-1}u^k(\varpi^{-2}t, \varpi^{-1}x), \quad k = 1, 2, \\ \delta a &= a^1 - a^2, \quad \delta v = v^1 - v^2. \end{aligned}$$

Thanks to (6.1), we find that $(\delta a, \delta v)$ satisfies

$$\begin{cases} \partial_t \delta a + v^2 \cdot \nabla \delta a = \delta F, \\ \partial_t \delta v - \mathcal{A} \delta v = \delta G, \\ (\delta a, \delta v)|_{t=0} = (0, 0), \end{cases} \quad (6.17)$$

where

$$\begin{aligned} \delta F &= -\delta v \cdot \nabla a^1 - \operatorname{div} \delta v - a^1 \operatorname{div} \delta v - \delta a \operatorname{div} v^2, \\ \delta G &= -\nabla \delta a - (v^1 \cdot \nabla v^1 - v^2 \cdot \nabla v^2) - L(a^1) \mathcal{A} v^1 + L(a^2) \mathcal{A} v^2 \\ &\quad - K(a^1) \nabla a^1 + K(a^2) \nabla a^2. \end{aligned}$$

In what follows, we denote $V^i(t) = \int_0^t \|v^i(\tau)\|_{\dot{B}_{p,1}^{\frac{n}{p}+1}} d\tau$ for $i = 1, 2$, and denote by A_T a constant depending on $\|a^1\|_{\tilde{L}_T^\infty(\dot{B}_{p,1}^{\frac{n}{p}})}$ and $\|a^2\|_{\tilde{L}_T^\infty(\dot{B}_{p,1}^{\frac{n}{p}})}$. Due to the inclusion relation $\mathcal{E}^{\frac{n}{p}} \subseteq \mathcal{E}^1$ ($p \leq n$), it suffices to prove the uniqueness of the solution in $\mathcal{E}^1$. So, in the following we take $p = n$.

Applying Proposition 3.4 gives

$$\|\delta a(t)\|_{\dot{B}_{p,\infty}^0} \leq e^{CV^2(t)} \int_0^t \|\delta F(\tau)\|_{\dot{B}_{p,\infty}^0} d\tau. \quad (6.18)$$

By Lemma 2.8, we have

$$\|\delta F(\tau)\|_{\dot{B}_{p,\infty}^0} \leq C\|v^2\|_{\dot{B}_{p,1}^2} \|\delta a\|_{\dot{B}_{p,\infty}^0} + C(1 + \|a^1\|_{\dot{B}_{p,1}^1}) \|\delta v\|_{\dot{B}_{p,1}^1}.$$

Plugging it into (6.18), we get by Gronwall's inequality that

$$\|\delta a(t)\|_{\dot{B}_{p,\infty}^0} \leq e^{CV^2(t)} \int_0^t (1 + \|a^1\|_{\dot{B}_{p,1}^1}) \|\delta v\|_{\dot{B}_{p,1}^1} d\tau. \quad (6.19)$$

Applying Proposition 3.3 to the second equation of (6.17) gives

$$\|\delta v(t)\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^1)} + \|\delta v(t)\|_{\tilde{L}_t^2(\dot{B}_{p,\infty}^0)} \leq C \int_0^t \|\delta G(\tau)\|_{\dot{B}_{p,\infty}^{-1}} d\tau. \quad (6.20)$$

From Lemma 2.8 and Proposition 7.5, we infer that

$$\begin{aligned} \|\delta G(t)\|_{\dot{B}_{p,\infty}^{-1}} &\leq C\|(v^1, v^2)\|_{\dot{B}_{p,1}^1} \|\delta v\|_{\dot{B}_{p,\infty}^0} + A_t \|a^1\|_{\dot{B}_{p,1}^1} \|\delta v\|_{\dot{B}_{p,\infty}^1} \\ &\quad + A_t(1 + \|v^2\|_{\dot{B}_{p,1}^2}) \|\delta a\|_{\dot{B}_{p,\infty}^0}. \end{aligned} \quad (6.21)$$

We take T small enough such that

$$\|(v^1, v^2)\|_{\tilde{L}_T^1(\dot{B}_{p,1}^2) \cap \tilde{L}_T^2(\dot{B}_{p,1}^1)} \ll 1.$$

Thus, plugging (6.21) into (6.20), we infer that for any $t \in [0, T]$,

$$\|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^1)} \leq A_T \int_0^t (1 + \|(v^1, v^2)\|_{\dot{B}_{p,1}^2}) \|\delta a\|_{\dot{B}_{p,\infty}^0} d\tau. \quad (6.22)$$

Lemma 6.5. [13] Let $s \in \mathbb{R}$. Then for any $1 \leq p, r \leq +\infty$ and $0 < \epsilon \leq 1$, we have

$$\|f\|_{\tilde{L}_T^r(\dot{B}_{p,1}^s)} \leq C \frac{\|f\|_{\tilde{L}_T^r(\dot{B}_{p,\infty}^s)}}{\epsilon} \log \left(e + \frac{\|f\|_{\tilde{L}_T^r(\dot{B}_{p,\infty}^{s-\epsilon})} + \|f\|_{\tilde{L}_T^r(\dot{B}_{p,\infty}^{s+\epsilon})}}{\|f\|_{\tilde{L}_T^r(\dot{B}_{p,\infty}^s)}} \right).$$

From Lemma 6.5, it follows that

$$\|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,1}^1)} \leq C \|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^1)} \log \left(e + \frac{\|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^0)} + \|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^2)}}{\|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^1)}} \right),$$

which together with (6.19) and (6.22) yields that for any $t \in [0, T]$,

$$\|\delta v\|_{\tilde{L}_t^1(\dot{B}_{p,\infty}^1)} \leq e^{CV^2(t)} A_T \int_0^t (1 + \|(v^1, v^2)\|_{\dot{B}_{p,1}^2}) \|\delta v\|_{\tilde{L}_\tau^1(\dot{B}_{p,\infty}^1)} \log(e + C_T \|\delta v\|_{\tilde{L}_\tau^1(\dot{B}_{p,\infty}^1)}^{-1}) d\tau,$$

where $C_T = \|\delta v\|_{\tilde{L}_T^1(\dot{B}_{p,\infty}^0)} + \|\delta v\|_{\tilde{L}_T^1(\dot{B}_{p,\infty}^2)}$. Noticing that $\|(v^1, v^2)(t)\|_{\dot{B}_{p,1}^2}$ is integrable on $[0, T]$ and

$$\int_0^1 \frac{dr}{r \log(r + C_T r^{-1})} dr = +\infty,$$

Osgood lemma applied concludes that $(\delta a, \delta v) = 0$ on $[0, T]$, and a continuity argument ensures that $(a^1, v^1) = (a^2, v^2)$ on $[0, \infty)$.

7. APPENDIX

In what follows, we denote $\chi_{\{\cdot\}}$ by the characteristic function defined in $\mathbb{Z}$, and $\{c(j)\}_{j \in \mathbb{Z}}$ by a sequence in ℓ^1 with the norm $\|\{c(j)\}\|_{\ell^1} \leq 1$.

Lemma 7.1. Let $s, \sigma, t, \tau \in \mathbb{R}$, $2 \leq p \leq 4$, p' is the conjugate index of p , and $1 \leq r, r_1, r_2 \leq \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$. Then there hold

(a) if $s \leq \frac{n}{2}$, $\sigma \leq \frac{n}{p}$, then for $2^j > R_0$,

$$\begin{aligned} \|\Delta_j(T_f g)\|_{L_T^r L^p} &\leq C c(j) \left(2^{j(\frac{n}{p'} - s - t)} + 2^{j(\frac{n}{2} - s - \tau)} + 2^{j(\frac{n}{p} - \sigma - \tau)} \right) \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}; \end{aligned} \quad (7.1)$$

(b) if $s \leq \frac{n}{p}$, $\sigma \leq \frac{2n}{p} - \frac{n}{2}$, then for $2^j \leq R_0$,

$$\begin{aligned} \|\Delta_j(T_f g)\|_{L_T^r L^2} &\leq C c(j) \left(2^{j(\frac{n}{2} - s - t)} + \chi_{\{2^j \sim R_0\}} \left(2^{j(\frac{n}{p} - s - \tau)} + 2^{j(\frac{2n}{p} - \frac{n}{2} - \sigma - \tau)} \right) \right) \\ &\quad \times \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}; \end{aligned} \quad (7.2)$$

(c) if $s \leq \frac{n}{2}$, $\sigma \leq \frac{n}{p}$, then

$$\|\Delta_j(T_f g)\|_{L_T^r L^p} \leq C c(j) \left(2^{j(\frac{n}{2}-s-t)} + 2^{j(\frac{n}{p}-\sigma-t)} \right) \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^t}. \quad (7.3)$$

Proof. (a) Thanks to (2.1), we have

$$\Delta_j(T_f g) = \sum_{|k-j| \leq 4} \Delta_j(S_{k-1} f \Delta_k g) = \sum_{|k-j| \leq 4} \sum_{k' \leq k-2} \Delta_j(\Delta_{k'} f \Delta_k g).$$

We denote $J \stackrel{\text{def}}{=} \{(k', k); |k-j| \leq 4, k' \leq k-2\}$, then for $2^j > R_0$,

$$\begin{aligned} \|\Delta_j(T_f g)\|_{L_T^r L^p} &\leq \sum_J \|\Delta_j(\Delta_{k'} f \Delta_k g)\|_{L_T^r L^p} \\ &\leq \left(\sum_{J_{\ell m}} + \sum_{J_{\ell h}} + \sum_{J_{hh}} \right) \|\Delta_j(\Delta_{k'} f \Delta_k g)\|_{L_T^r L^p} \\ &\stackrel{\text{def}}{=} I_1 + I_2 + I_3, \end{aligned}$$

where

$$\begin{aligned} J_{\ell m} &= \{(k', k) \in J, 2^{k'} \leq R_0, 2^{-4} R_0 \leq 2^k \leq R_0\}, \\ J_{\ell h} &= \{(k', k) \in J, 2^{k'} \leq R_0, 2^k > R_0\}, \\ J_{hh} &= \{(k', k) \in J, 2^{k'} > R_0, 2^k > R_0\}. \end{aligned}$$

We get by using Lemma 2.1 and $s \leq \frac{n}{2}$ that

$$\begin{aligned} I_1 &\leq C \sum_{(k', k) \in J_{\ell m}} \|\Delta_{k'} f\|_{L_T^{r_1} L^\infty} 2^{k(\frac{n}{2}-\frac{n}{p})} \|\Delta_k g\|_{L_T^{r_2} L^2} \\ &\leq C \sum_{(k', k) \in J_{\ell m}} 2^{k's} \|\Delta_{k'} f\|_{L_T^{r_1} L^2} 2^{k'(\frac{n}{2}-s)} 2^{kt} \|\Delta_k g\|_{L_T^{r_2} L^2} 2^{k(\frac{n}{2}-\frac{n}{p}-t)} \\ &\leq C c(j) 2^{j(\frac{n}{p}-s-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} I_2 &\leq \sum_{(k', k) \in J_{\ell h}} \|\Delta_{k'} f\|_{L_T^{r_1} L^\infty} \|\Delta_k g\|_{L_T^{r_2} L^p} \\ &\leq C \sum_{(k', k) \in J_{\ell h}} 2^{k's} \|\Delta_{k'} f\|_{L_T^{r_1} L^2} 2^{k'(\frac{n}{2}-s)} 2^{k\tau} \|\Delta_k g\|_{L_T^{r_2} L^p} 2^{-k\tau} \\ &\leq C c(j) 2^{j(\frac{n}{2}-s-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and noting $\sigma \leq \frac{n}{p}$,

$$\begin{aligned} I_3 &\leq \sum_{(k', k) \in J_{hh}} \|\Delta_{k'} f\|_{L_T^{r_1} L^\infty} \|\Delta_k g\|_{L_T^{r_2} L^p} \\ &\leq C \sum_{(k', k) \in J_{hh}} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^{r_1} L^p} 2^{k'(\frac{n}{p}-\sigma)} 2^{k\tau} \|\Delta_k g\|_{L_T^{r_2} L^p} 2^{-k\tau} \\ &\leq C c(j) 2^{j(\frac{n}{p}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Then the inequality (7.1) can be deduced from the estimates of I_1 , I_2 and I_3 .

(b) We denote

$$J \stackrel{\text{def}}{=} \{(k', k); |k - j| \leq 4, k' \leq k - 2\},$$

then for $2^j \leq R_0$, there holds

$$J = J_{\ell\ell} \cup J_{\ell m} \cup J_{hm},$$

where

$$\begin{aligned} J_{\ell\ell} &= \{(k', k) \in J, 2^{k'} \leq R_0, 2^k \leq R_0\}, \\ J_{\ell m} &= \{(k', k) \in J, 2^{k'} \leq R_0, R_0 < 2^k \leq 2^4 R_0\}, \\ J_{hm} &= \{(k', k) \in J, 2^{k'} > R_0, R_0 < 2^k \leq 2^4 R_0\}. \end{aligned}$$

Then thanks to (2.1), we have for $2^j \leq R_0$,

$$\begin{aligned} \|\Delta_j(Tfg)\|_{L_T^r L^2} &\leq \sum_J \|\Delta_j(\Delta_{k'} f \Delta_k g)\|_{L_T^r L^2} \\ &\leq \left(\sum_{J_{\ell\ell}} + \sum_{J_{\ell m}} + \sum_{J_{hm}} \right) \|\Delta_j(\Delta_{k'} f \Delta_k g)\|_{L_T^r L^2} \\ &\stackrel{\text{def}}{=} II_1 + II_2 + II_3. \end{aligned}$$

We get by using Lemma 2.1 and $s \leq \frac{n}{p}$ that

$$\begin{aligned} II_1 &\leq C \sum_{(k', k) \in J_{\ell\ell}} 2^{k's} \|\Delta_{k'} f\|_{L_T^{r_1} L^2} 2^{k'(\frac{n}{2}-s)} 2^{kt} \|\Delta_k g\|_{L_T^{r_2} L^2} 2^{-kt} \\ &\leq Cc(j) 2^{j(\frac{n}{2}-s-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \\ II_2 &\leq C \sum_{(k', k) \in J_{\ell m}} 2^{k's} \|\Delta_{k'} f\|_{L_T^{r_1} L^2} 2^{k'(\frac{n}{p}-s)} 2^{k\tau} \|\Delta_k g\|_{L_T^{r_2} L^p} 2^{-k\tau} \\ &\leq Cc(j) \chi_{\{2^j \sim R_0\}} 2^{j(\frac{n}{p}-s-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and noting $\sigma \leq \frac{2n}{p} - \frac{n}{2}$,

$$\begin{aligned} II_3 &\leq C \sum_{(k', k) \in J_{hm}} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^{r_1} L^p} 2^{k'(\frac{2n}{p}-\frac{n}{2}-\sigma)} 2^{k\tau} \|\Delta_k g\|_{L_T^{r_2} L^p} 2^{-k\tau} \\ &\leq Cc(j) \chi_{\{2^j \sim R_0\}} 2^{j(\frac{2n}{p}-\frac{n}{2}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

The inequality (7.2) follows from the above estimates.

(c) Since the proof of (7.3) is similar, here we omit it. $\square$

Lemma 7.2. Let $s, \sigma, t, \tau \in \mathbb{R}$, $2 \leq p \leq 4$, p' is the conjugate index of p , and $1 \leq r, r_1, r_2 \leq \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$. Assume that $s + t > 0$, $s + \tau > 0$, $\sigma + t > 0$, and

$\sigma + \tau > 0$. Then there hold

$$\begin{aligned} \|\Delta_j R(f, g)\|_{L_T^r L^p} &\leq Cc(j) \left(2^{j(\frac{n}{p'} - s - t)} + 2^{j(\frac{n}{2} - s - \tau)} \right. \\ &\quad \left. + 2^{j(\frac{n}{2} - \sigma - t)} + 2^{j(\frac{n}{p} - \sigma - \tau)} \right) \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}; \end{aligned} \quad (7.4)$$

$$\begin{aligned} \|\Delta_j R(f, g)\|_{L_T^r L^2} &\leq Cc(j) \left(2^{j(\frac{n}{2} - s - t)} + 2^{j(\frac{n}{p} - s - \tau)} \right. \\ &\quad \left. + 2^{j(\frac{n}{p} - \sigma - t)} + 2^{j(\frac{2n}{p} - \frac{n}{2} - \sigma - \tau)} \right) \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}; \end{aligned} \quad (7.5)$$

$$\|\Delta_j R(f, g)\|_{L_T^r L^2} \leq Cc(j) \left(2^{j(\frac{n}{2} - s - t)} + 2^{j(\frac{n}{p} - s - \tau)} \right) \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,1}^s} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \quad (7.6)$$

where p' is the conjugate index of p .

Proof. Thanks to (2.1), we have

$$\Delta_j(R(f, g)) = \sum_{k \geq j-3} \sum_{|k' - k| \leq 1} \Delta_j(\Delta_k f \Delta_{k'} g),$$

Set $J \stackrel{\text{def}}{=} \{(k, k'); k \geq j-3, |k' - k| \leq 1\}$, then

$$J = J_{\ell\ell} \cup J_{\ell m} \cup J_{hm} \cup J_{hh},$$

where

$$\begin{aligned} J_{\ell\ell} &= \{(k, k') \in J, 2^k \leq R_0, 2^{k'} \leq R_0\}, \\ J_{\ell m} &= \{(k, k') \in J, 2^k \leq R_0, R_0 < 2^{k'} \leq 2R_0\}, \\ J_{hm} &= \{(k, k') \in J, 2^k > R_0, 2^{-1}R_0 \leq 2^{k'} \leq R_0\}, \\ J_{hh} &= \{(k, k') \in J, 2^k > R_0, 2^{k'} > R_0\}. \end{aligned}$$

Thus, we have

$$\begin{aligned} \Delta_j(R(f, g)) &= \left(\sum_{J_{\ell\ell}} + \sum_{J_{\ell m}} + \sum_{J_{hm}} + \sum_{J_{hh}} \right) \Delta_j(\Delta_k f \Delta_{k'} g) \\ &\stackrel{\text{def}}{=} I_1 + I_2 + I_3 + I_4. \end{aligned}$$

We get by Lemma 2.1 and $s + t > 0$ that

$$\begin{aligned} \|I_1\|_{L_T^r L^p} &\leq C 2^{j\frac{n}{p'}} \sum_{(k, k') \in J_{\ell\ell}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^1} \\ &\leq C 2^{j\frac{n}{p'}} \sum_{(k, k') \in J_{\ell\ell}} 2^{ks} \|\Delta_k f\|_{L_T^{r_1} L^2} 2^{-ks} 2^{k't} \|\Delta_{k'} g\|_{L_T^{r_2} L^2} 2^{-k't} \\ &\leq Cc(j) 2^{j(\frac{n}{p'} - s - t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and

$$\begin{aligned} \|I_1\|_{L_T^r L^2} &\leq C 2^{j\frac{n}{2}} \sum_{(k, k') \in J_{\ell\ell}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^1} \\ &\leq Cc(j) 2^{j(\frac{n}{2} - s - t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Similarly, due to $s + \tau > 0$, we obtain

$$\begin{aligned} \|I_2\|_{L_T^r L^p} &\leq C 2^{j\frac{n}{2}} \sum_{(k,k') \in J_{\ell m}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{2p}{2+p}}} \\ &\leq C 2^{j\frac{n}{2}} \sum_{(k,k') \in J_{\ell m}} 2^{ks} \|\Delta_k f\|_{L_T^{r_1} L^2} 2^{-ks} 2^{k'\tau} \|\Delta_{k'} g\|_{L_T^{r_2} L^p} 2^{-k'\tau} \\ &\leq C c(j) 2^{j(\frac{n}{2}-s-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and

$$\begin{aligned} \|I_2\|_{L_T^r L^2} &\leq C 2^{j\frac{n}{p}} \sum_{(k,k') \in J_{\ell m}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{2p}{2+p}}} \\ &\leq C c(j) 2^{j(\frac{n}{p}-s-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Thanks to $\sigma + t > 0$, we have

$$\begin{aligned} \|I_3\|_{L_T^r L^p} &\leq C 2^{j\frac{n}{2}} \sum_{(k,k') \in J_{hm}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{2p}{2+p}}} \\ &\leq C 2^{j\frac{n}{2}} \sum_{(k,k') \in J_{hm}} 2^{k\sigma} \|\Delta_k f\|_{L_T^{r_1} L^p} 2^{-k\sigma} 2^{k't} \|\Delta_{k'} g\|_{L_T^{r_2} L^2} 2^{-k't} \\ &\leq C c(j) 2^{j(\frac{n}{2}-\sigma-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and

$$\begin{aligned} \|I_3\|_{L_T^r L^2} &\leq C 2^{j\frac{n}{p}} \sum_{(k,k') \in J_{hm}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{2p}{2+p}}} \\ &\leq C c(j) 2^{j(\frac{n}{p}-\sigma-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Finally, due to $\sigma + \tau > 0$ and $2 \leq p \leq 4$, we have

$$\begin{aligned} \|I_4\|_{L_T^r L^p} &\leq C 2^{j\frac{n}{p}} \sum_{(k,k') \in J_{hh}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{p}{2}}} \\ &\leq C 2^{j\frac{n}{p}} \sum_{(k,k') \in J_{hh}} 2^{k\sigma} \|\Delta_k f\|_{L_T^{r_1} L^p} 2^{-k\sigma} 2^{k'\tau} \|\Delta_{k'} g\|_{L_T^{r_2} L^p} 2^{-k'\tau} \\ &\leq C c(j) 2^{j(\frac{n}{p}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and

$$\begin{aligned} \|I_4\|_{L_T^r L^2} &\leq C 2^{j(\frac{2n}{p}-\frac{n}{2})} \sum_{(k,k') \in J_{hh}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{p}{2}}} \\ &\leq C c(j) 2^{j(\frac{2n}{p}-\frac{n}{2}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

Then the inequalities (7.4) and (7.5) can be deduced from the above estimates. Finally, the inequality (7.6) can be deduced from

$$\begin{aligned} \|I_1 + I_3\|_{L_T^r L^2} &\leq C 2^{j\frac{n}{2}} \sum_{(k,k') \in J_{\ell\ell} \cup J_{hm}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^1} \\ &\leq C c(j) 2^{j(\frac{n}{2}-s-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,1}^s} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}, \end{aligned}$$

and

$$\begin{aligned} \|I_2 + I_4\|_{L_T^r L^2} &\leq C 2^{j\frac{n}{p}} \sum_{(k,k') \in J_{\ell m} \cup J_{hh}} \|\Delta_k f \Delta_{k'} g\|_{L_T^r L^{\frac{2p}{2+p}}} \\ &\leq C c(j) 2^{j(\frac{n}{p}-s-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,1}^s} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,\tau}}. \end{aligned}$$

This completes the proof of Lemma 7.2. $\square$

Proposition 7.3. Let $s, t, \tilde{s}, \tilde{t}, \sigma, \tau \in \mathbb{R}$, $2 \leq p \leq 4$, and $1 \leq r, r_1, r_2 \leq \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$. Then we have

(a) if $\sigma, \tau \leq \frac{n}{p}$ and $\sigma + \tau > 0$, then

$$\sum_{2^j > R_0} 2^{j(\sigma+\tau-\frac{n}{p})} \|\Delta_j(fg)\|_{L_T^r L^p} \leq C \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\tau}}; \quad (7.7)$$

(b) if $s, \tilde{s} \leq \frac{n}{p}$, $s + t > n - \frac{2n}{p}$ with $s + t = \tilde{s} + \tilde{t}$, and $\gamma \in \mathbb{R}$, then

$$\begin{aligned} &\sum_{2^j \leq R_0} 2^{j(s+t-\frac{n}{2})} \|\Delta_j(fg)\|_{L_T^r L^2} \\ &\leq C \left(\|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,s-\frac{n}{2}+\frac{n}{p}}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,t-\frac{n}{2}+\frac{n}{p}+\gamma}} + \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\tilde{s},\tilde{s}-\frac{n}{2}+\frac{n}{p}}} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\tilde{t},\tilde{t}-\frac{n}{2}+\frac{n}{p}}} \right). \end{aligned} \quad (7.8)$$

(c) if $s, \tilde{s} \leq \frac{n}{2}$, $s + t > \frac{n}{2} - \frac{n}{p}$ with $s + t = \tilde{s} + \tilde{t}$, then

$$\begin{aligned} &\sum_{j \in \mathbb{Z}} 2^{j(s+t-\frac{n}{2})} \|\Delta_j(fg)\|_{L_T^r L^2} \\ &\leq C \left(\|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{s,s-\frac{n}{2}+\frac{n}{p}}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^{t,t-\frac{n}{2}+\frac{n}{p}}} + \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\tilde{s},\tilde{s}-\frac{n}{2}+\frac{n}{p}}} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,1}^{\tilde{t},\tilde{t}-\frac{n}{2}+\frac{n}{p}}} \right). \end{aligned} \quad (7.9)$$

Proof. Thanks to (7.1), we get for $2^j > R_0$,

$$\|\Delta_j(T_f g)\|_{L_T^r L^p} + \|\Delta_j(T_g f)\|_{L_T^r L^p} \leq C c(j) 2^{j(\frac{n}{p}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\tau}},$$

and from (7.4), we infer that

$$\|\Delta_j(R(f, g))\|_{L_T^r L^p} \leq C c(j) 2^{j(\frac{n}{p}-\sigma-\tau)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\sigma}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\frac{n}{2}-\frac{n}{p}+\tau}},$$

from which and Bony' decomposition (2.2), we obtain (7.7).

From (7.2), we get for $2^j \leq R_0$,

$$\begin{aligned} \|\Delta_j(T_f g)\|_{L_T^r L^2} &\leq C c(j) 2^{j(\frac{n}{2}-s-t)} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s,s-\frac{n}{2}+\frac{n}{p}}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{t,t-\frac{n}{2}+\frac{n}{p}+\gamma}}, \\ \|\Delta_j(T_g f)\|_{L_T^r L^2} &\leq C c(j) 2^{j(\frac{n}{2}-s-t)} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\tilde{s},\tilde{s}-\frac{n}{2}+\frac{n}{p}}} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\tilde{t},\tilde{t}-\frac{n}{2}+\frac{n}{p}}}, \end{aligned}$$

and from (7.5),

$$\|\Delta_j(R(f, g))\|_{L_T^r L^2} \leq C c(j) 2^{j(\frac{n}{2}-s-t)} \|f\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\tilde{t},\tilde{t}-\frac{n}{2}+\frac{n}{p}}} \|g\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{\tilde{s},\tilde{s}-\frac{n}{2}+\frac{n}{p}}},$$

which imply (7.8). In the same manner, the inequality (7.9) can be deduced from (7.3) and (7.6). $\square$

Proposition 7.4. Let $2 \leq p \leq 4$, $-\frac{n}{p} < s \leq \frac{n}{2} + 1$, and $-\frac{n}{p} < \sigma \leq \frac{n}{p} + 1$, and $1 \leq r, r_1, r_2 \leq \infty$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$. Then for $2^j \geq R_0$, there holds

$$\|[v, \Delta_j] \cdot \nabla f\|_{L_T^r L^p} \leq Cc(j)(2^{-j\sigma} + 2^{j(\frac{n}{2} - \frac{n}{p} - s)})\|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}\|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s, \sigma}}. \quad (7.10)$$

Moreover, if $-\frac{n}{p} < s \leq \frac{n}{p} + 1$, then

$$\|[v, \Delta_j] \cdot \nabla f\|_{L_T^r L^2} \leq Cc(j)2^{-js}\|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}\|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^s}. \quad (7.11)$$

Proof. Using the Bony's decomposition (2.2), we write

$$\begin{aligned} [v, \Delta_j] \cdot \nabla f &= [T_{v^i}, \Delta_j] \partial_i f + T_{\partial_i \Delta_j f} v^i + R(v^i, \partial_i \Delta_j f) \\ &\quad - \Delta_j(T_{\partial_i f} v^i) - \Delta_j R(v^i, \partial_i f). \end{aligned}$$

We get by using (7.1), $s \leq \frac{n}{2} + 1$ and $\sigma \leq \frac{n}{p} + 1$ that

$$\|\Delta_j(T_{\partial_i f} v^i)\|_{L_T^r L^p} \leq Cc(j)(2^{-j\sigma} + 2^{j(\frac{n}{2} - \frac{n}{p} - s)})\|\nabla f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s-1, \sigma-1}}\|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}.$$

And from (7.4), $s > -\frac{n}{p}$ and $\sigma > -\frac{n}{p}$, we infer that

$$\|\Delta_j R(v^i, \partial_i f)\|_{L_T^r L^p} \leq Cc(j)(2^{-j\sigma} + 2^{j(\frac{n}{2} - \frac{n}{p} - s)})\|\nabla f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s-1, \sigma-1}}\|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}.$$

Noticing that

$$T'_{\partial_i \Delta_j f} v^i \stackrel{\text{def}}{=} T_{\partial_i \Delta_j f} v^i + R(v^i, \partial_i \Delta_j f) = \sum_{k \geq j-2} S_{k+2} \Delta_j \partial_i f \Delta_k v^i,$$

then we get by Lemma 2.1 that

$$\begin{aligned} \|T'_{\partial_i \Delta_j f} v^i\|_{L_T^r L^p} &\leq C\|\Delta_j \nabla f\|_{L_T^{r_2} L^\infty} \sum_{k \geq j-2} \|\Delta_k v\|_{L_T^{r_1} L^p} \\ &\leq C2^{j(1+\frac{n}{p})} \|\Delta_j f\|_{L_T^{r_2} L^p} \left(\sum_{k \geq j-2, 2^k > R_0} \|\Delta_k v\|_{L_T^{r_1} L^p} + \sum_{k \geq j-2, \frac{R_0}{4} \leq 2^k \leq R_0} 2^{k(\frac{n}{2} - \frac{n}{p})} \|\Delta_k v\|_{L_T^{r_1} L^2} \right) \\ &\leq Cc(j)2^{-j\sigma} \|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s, \sigma}}. \end{aligned} \quad (7.12)$$

Now we turn to estimate $[T_{v^i}, \Delta_j] \partial_i f = \sum_{|k-j| \leq 4} [S_{k-1} v^i, \Delta_j] \partial_i \Delta_k f$. Set $h(x) = (\mathcal{F}^{-1} \phi)(x)$, we get by integration by parts that

$$\begin{aligned} [T_{v^i}, \Delta_j] \partial_i f &= \sum_{|k-j| \leq 4} 2^{nj} \int_{\mathbb{R}^n} h(2^j(x-y))(S_{k-1} v^i(x) - S_{k-1} v^i(y)) \partial_i \Delta_k f(y) dy \\ &= \sum_{|k-j| \leq 4} 2^{(n+1)j} \int_{\mathbb{R}^n} \int_0^1 y \cdot \nabla S_{k-1} v^i(x - \tau y) d\tau \partial_i h(2^j y) \Delta_k f(x-y) dy \\ &\quad + 2^{nj} \int_{\mathbb{R}^n} h(2^j(x-y)) \partial_i S_{k-1} v^i(y) \Delta_k f(y) dy, \end{aligned}$$

which together with the Minkowski inequality and $\dot{B}_{2,p}^{\frac{n}{2}, \frac{n}{p}} \hookrightarrow L^\infty$ implies

$$\begin{aligned} & \| [T_{v^i}, \Delta_j] \partial_i f \|_{L_T^r L^p} \\ & \leq C \|\nabla v\|_{L_T^{r_1} L^\infty} \left(\sum_{|k-j| \leq 4, 2^k > R_0} \|\Delta_k f\|_{L_T^{r_2} L^p} + \sum_{|k-j| \leq 4, \frac{R_0}{2^4} \leq 2^k \leq R_0} 2^{k(\frac{n}{2} - \frac{n}{p})} \|\Delta_k f\|_{L_T^{r_2} L^2} \right) \\ & \leq C c(j) (2^{-j\sigma} + 2^{j(\frac{n}{2} - \frac{n}{p} - s)}) \|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,p}^{s, \sigma}}. \end{aligned}$$

Summing up the above estimates, we conclude the inequality (7.10). On the other hand, we have from (7.6),

$$\|\Delta_j R(v^i, \partial_i f)\|_{L_T^r L^2} \leq C c(j) 2^{-js} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^s} \|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}}.$$

With a slightly modification of (7.12), we get

$$\|\Delta_j (T_{\partial_i f} v^i)\|_{L_T^r L^2} + \|T'_{\partial_i \Delta_j f} v^i\|_{L_T^r L^2} \leq C c(j) 2^{-js} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^s} \|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}},$$

and thanks to the representation of $[T_{v^i}, \Delta_j] \partial_i f$, we have

$$\begin{aligned} \|[T_{v^i}, \Delta_j] \partial_i f\|_{L_T^r L^2} & \leq C \|\nabla v\|_{L_T^{r_1} L^\infty} \sum_{|k-j| \leq 4} \|\Delta_k f\|_{L_T^{r_2} L^2} \\ & \leq C c(j) 2^{-js} \|v\|_{\tilde{L}_T^{r_1} \dot{B}_{2,p}^{\frac{n}{2}+1, \frac{n}{p}+1}} \|f\|_{\tilde{L}_T^{r_2} \dot{B}_{2,1}^s}. \end{aligned}$$

Then the inequality (7.11) follows from the above three estimates. $\square$

Proposition 7.5. Let $2 \leq p \leq 4$, $s, \sigma > 0$, and $s \geq \sigma - \frac{n}{2} + \frac{n}{p}$, $r \geq 1$. Assume that $F \in W_{\text{loc}}^{[s]+2, \infty} \cap W_{\text{loc}}^{[\sigma]+2, \infty}$ with $F(0) = 0$. Then there holds

$$\|F(f)\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s, \sigma}} \leq C (1 + \|f\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}})^{\max([s], [\sigma])+1} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s, \sigma}}. \quad (7.13)$$

For any $s > 0$ and $p \geq 1$, there holds

$$\|F(f)\|_{\tilde{L}_T^r \dot{B}_{p,1}^s} \leq C (1 + \|f\|_{L_T^\infty L^\infty})^{[s]+1} \|f\|_{\tilde{L}_T^r \dot{B}_{p,1}^s}. \quad (7.14)$$

Proof. The inequality (7.14) is classical, see [12]. We only present the proof of (7.13). Decompose $F(f)$ as

$$\begin{aligned} F(f) &= \sum_{k' \in \mathbb{Z}} F(S_{k'+1} f) - F(S_{k'} f) = \sum_{k' \in \mathbb{Z}} \Delta_{k'} f \int_0^1 F'(S_{k'} f + \tau \Delta_{k'} f) d\tau \\ &\stackrel{\text{def}}{=} \sum_{k' \in \mathbb{Z}} \Delta_{k'} f m_{k'}, \end{aligned}$$

where $m_{k'} \stackrel{\text{def}}{=} \int_0^1 F'(S_{k'} f + \tau \Delta_{k'} f) d\tau$. We denote

$$J_\ell = \{k; 2^k \leq R_0\}, \quad J_h = \{k; 2^k > R_0\}.$$

Then we have

$$J_\ell = J_{\ell\ell} \cup J_{\ell m} \cup J_{\ell h}, \quad J_h = J_{h\ell} \cup J_{hm} \cup J_{hh}.$$

where

$$\begin{aligned}
J_{\ell\ell} &= \{(k, k'); k \in J_\ell, k' \leq k\}, \\
J_{\ell m} &= \{(k, k'); k \in J_\ell, k' > k, 2^{k'} \leq R_0\}, \\
J_{\ell h} &= \{(k, k'); k \in J_\ell, k' > k, 2^{k'} > R_0\}, \\
J_{h\ell} &= \{(k, k'); k \in J_h, k' \leq k, 2^{k'} \leq R_0\}, \\
J_{hm} &= \{(k, k'); k \in J_h, k' \leq k, 2^{k'} > R_0\}, \\
J_{hh} &= \{(k, k'); k \in J_h, k' > k\}.
\end{aligned}$$

By Lemma 2.1, we have

$$\|\Delta_k(\Delta_{k'} f m_{k'})\|_{L^2} \leq C 2^{-k|\alpha|} \sup_{|\gamma|=|\alpha|} \|D^\gamma \Delta_k(\Delta_{k'} f m_{k'})\|_{L^2},$$

for any $\alpha \in \mathbb{N}^n$ and for $|\gamma| \geq 0$,

$$\|D^\gamma m_{k'}\|_{L^\infty} \leq C 2^{k'|\gamma|} (1 + \|f\|_{L^\infty})^{|\gamma|},$$

which imply

$$\|\Delta_k(\Delta_{k'} f m_{k'})\|_{L^2} \leq C 2^{(k'-k)|\alpha|} \|\Delta_{k'} f\|_{L^2} (1 + \|f\|_{L^\infty})^{|\alpha|}. \quad (7.15)$$

We apply (7.15) with $|\alpha| = [s] + 1$ to get

$$\begin{aligned}
& \sum_{(k,k') \in J_{\ell\ell}} 2^{ks} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^2} \\
& \leq C \sum_{2^{k'} \leq R_0} 2^{k's} \|\Delta_{k'} f\|_{L^2} \sum_{k \geq k'} 2^{(k-k')(s-[s]-1)} (1 + \|f\|_{L_T^\infty L^\infty})^{|\alpha|} \\
& \leq C (1 + \|f\|_{L^\infty})^{[s]+1} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}.
\end{aligned} \quad (7.16)$$

And we apply (7.15) with $|\alpha| = 0$ to get

$$\begin{aligned}
\sum_{(k,k') \in J_{\ell m}} 2^{ks} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^2} & \leq C \sum_{2^{k'} \leq R_0} 2^{k's} \|\Delta_{k'} f\|_{L_T^r L^2} \sum_{k < k'} 2^{(k-k')s} \\
& \leq C \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}.
\end{aligned} \quad (7.17)$$

Setting $m_0 \stackrel{\text{def}}{=} F'(0)$, we write

$$\Delta_k(\Delta_{k'} f m_{k'}) = \Delta_k(\Delta_{k'} f m_0) + \Delta_k(\Delta_{k'} f (m_{k'} - m_0)).$$

It is easy to find that

$$\sum_{(k,k') \in J_{\ell h}} 2^{ks} \|\Delta_k(\Delta_{k'} f m_0)\|_{L_T^r L^2} \leq C \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}},$$

and using the formula

$$\begin{aligned}
m_{k'} - m_0 &= \int_0^1 m'_{k'}(\tau(S_{k'-1}f + \Delta_{k'}f)) d\tau(S_{k'-1}f + \Delta_{k'}f) \\
&\stackrel{\text{def}}{=} (S_{k'-1}f + \Delta_{k'}f) \tilde{m}_{k'},
\end{aligned}$$

we obtain

$$\begin{aligned}
& \sum_{(k,k') \in J_{\ell h}} 2^{ks} \|\Delta_k(\Delta_{k'} f(m_{k'} - m_0))\|_{L_T^r L^2} \\
& \leq C \sum_{(k,k') \in J_{\ell h}} 2^{ks} (\|\Delta_k(\Delta_{k'} f \tilde{m}_{k'} S_{k'-1} f)\|_{L_T^r L^2} + 2^{k(\frac{2n}{p} - \frac{n}{2})} \|(\Delta_{k'} f)^2\|_{L_T^r L^{\frac{p}{2}}}) \\
& \stackrel{\text{def}}{=} I_1 + I_2.
\end{aligned}$$

Now we have

$$\begin{aligned}
I_1 & \leq C \sum_{(k,k') \in J_{\ell h}} 2^{ks} \|\Delta_{k'} f\|_{L_T^r L^p} \sum_{k'' \leq k'-2, 2^{k''} \leq R_0} \|\Delta_{k''} f\|_{L_T^\infty L^{\frac{2p}{p-2}}} \\
& + C \sum_{(k,k') \in J_{\ell h}} 2^{ks} 2^{k(\frac{2n}{p} - \frac{n}{2})} \sum_{k'' \leq k'-2, 2^{k''} > R_0} \|\Delta_{k'} f \Delta_{k''} f\|_{L_T^r L^{\frac{p}{2}}}.
\end{aligned}$$

The first term of the right hand side is bounded by

$$\begin{aligned}
& \sum_{2^{k'} \geq R_0} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^r L^p} 2^{-k'\sigma} \sum_{2^k \leq R_0} 2^{ks} \sum_{2^{k''} \leq R_0} 2^{k'' \frac{n}{p}} \|\Delta_{k''} f\|_{L_T^\infty L^2} \\
& \leq C \|f\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}},
\end{aligned}$$

and the second term is bounded by

$$\begin{aligned}
& \sum_{J_{\ell h}} 2^{k(s + \frac{2n}{p} - \frac{n}{2})} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^r L^p} 2^{-k'\sigma} \sum_{2^{k''} > R_0} 2^{k'' \frac{n}{p}} \|\Delta_{k''} f\|_{L_T^\infty L^p} 2^{-k'' \frac{n}{p}} \\
& \leq C \|f\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}.
\end{aligned}$$

Similarly, we have

$$I_2 \leq C \|f\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}.$$

Thus we obtain

$$\sum_{(k,k') \in J_{\ell h}} 2^{ks} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^2} \leq C(1 + \|f\|_{\tilde{L}_T^\infty \dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}}}) \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}. \quad (7.18)$$

Using the argument as leading to (7.16) and $s \geq \sigma - \frac{n}{2} + \frac{n}{p}$, we have

$$\begin{aligned}
& \sum_{(k,k') \in J_{h\ell}} 2^{k\sigma} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^p} \\
& \leq C(1 + \|f\|_{L_T^\infty L^\infty})^{[\sigma]+1} \sum_{2^{k'} \leq R_0} 2^{k's} \|\Delta_{k'} f\|_{L_T^r L^2} \sum_{k \geq k'} 2^{(k-k')(\sigma - [\sigma] - 1)} 2^{k'(\frac{n}{2} - \frac{n}{p} - s + \sigma)} \\
& \leq C(1 + \|f\|_{L_T^\infty L^\infty})^{[\sigma]+1} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}},
\end{aligned} \quad (7.19)$$

and

$$\begin{aligned}
& \sum_{(k,k') \in J_{hm}} 2^{k\sigma} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^p} \\
& \leq C(1 + \|f\|_{L_T^\infty L^\infty})^{[\sigma]+1} \sum_{2^{k'} > R_0} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^r L^p} \sum_{k > k'} 2^{(k-k')(\sigma-[\sigma]-1)} \\
& \leq C(1 + \|f\|_{L_T^\infty L^\infty})^{[\sigma]+1} \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}. \tag{7.20}
\end{aligned}$$

Finally due to $\sigma > 0$, we have

$$\begin{aligned}
\sum_{(k,k') \in J_{hh}} 2^{k\sigma} \|\Delta_k(\Delta_{k'} f m_{k'})\|_{L_T^r L^p} & \leq C \sum_{2^{k'} \geq R_0} 2^{k'\sigma} \|\Delta_{k'} f\|_{L_T^r L^p} \sum_{k' \geq k} 2^{(k-k')\sigma} \\
& \leq C \|f\|_{\tilde{L}_T^r \dot{B}_{2,p}^{s,\sigma}}. \tag{7.21}
\end{aligned}$$

Summing up (7.16)–(7.21) and noting $\dot{B}_{2,p}^{\frac{n}{p}, \frac{n}{p}} \hookrightarrow L^\infty$, we obtain (7.13). $\square$

8. ACKNOWLEDGMENTS

Q. Chen and C. Miao were partially supported by the NSF of China (grants 10701012 and 10725102). Z. Zhang was supported by the NSF of China (grant 10601002).

REFERENCES

- [1] H. Abidi, *Équation de Navier-Stokes avec densité et viscosité variables dans l'espace critique*. Rev. Mat. Iberoam. 23(2007), 537–586.
- [2] J.-M. Bony, *Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires*. Ann. de l'Ecole Norm. Sup., 14(1981), 209–246.
- [3] M. Cannone, *A generalization of a theorem by Kato on Navier-Stokes equations*. Revista Matemática Iberoamericana, 13(1997)515–541.
- [4] M. Cannone, Y. Meyer and F. Planchon, *Solutions autosimilaires des équations de Navier-Stokes*. Séminaire “Équations aux Dérivées Partielles” de l'École polytechnique, Exposé VIII, 1993–1994.
- [5] M. Cannone, C. Miao and G. Wu, *On the inviscid limit of the two-dimensional Navier-Stokes equations with fractional diffusion*. Advances in Mathematical Science and Application, 18(2008), 607–624.
- [6] J.-Y. Chemin, *Perfect incompressible fluids*. Oxford University Press, New York, 1998.
- [7] J.-Y. Chemin, *Localization in Fourier space and Navier-Stokes system*. Phase space analysis of partial differential equations. Vol. I, 2004, 53–135.
- [8] J.-Y. Chemin and I. Gallagher, *On the global wellposedness of the 3-D Navier-Stokes equations with large initial data*. Ann. de l'Ecole Norm. Sup., (39)2006, 679–698.
- [9] J.-Y. Chemin, I. Gallagher and M. Paicu, *Global regularity for some classes of large solutions to the Navier-Stokes equations*. preprint, 2008.
- [10] J.-Y. Chemin and P. Zhang, *On the global wellposedness of the 3-D incompressible anisotropic Navier-Stokes equations*. Comm. Math. Phys., (272)2007, 529–566.
- [11] R. Danchin, *Global existence in critical spaces for compressible Navier-Stokes equations*. Invent. Math., 141(2000), 579–614.
- [12] R. Danchin, *Local theory in critical spaces for compressible viscous and heat-conductive gases*. Comm. Partial Differential Equations, 26(2001), 1183–1233.
- [13] R. Danchin, *Density-dependent incompressible viscous fluids in critical spaces*. Proc. Roy. Soc. Edinburgh Sect. A, 133(2003), 1311–1334.
- [14] R. Danchin, *On the uniqueness in critical spaces for compressible Navier-Stokes equations*. Non-linear Differential Equations Appl., 12(2005), 111–128.
- [15] R. Danchin, *Well-posedness in critical spaces for barotropic viscous fluids with truly not constant density*. Comm. Partial Differential Equations, 32(2007), 1–17.

- [16] R. Danchin, *Uniform estimates for Transport-Diffusion equations*. J. Hyper. Differential Equations, 32(2007), 1373-1397.
- [17] H. Fujita and T. Kato, *On the Navier-Stokes initial value problem I*. Arch. Rational Mech. Anal., 16(1964), 269-315.
- [18] T. Hmidi and S. Keraani, *Global solutions of the super-critical 2D Q-G equation in Besov spaces*, Advances in Math., 214(2007), 618-638.
- [19] T. Hmidi and S. Keraani, *Incompressible viscous flows in borderline Besov space*. Arch. Rational Mech. Anal., 189(2008), 283-300.
- [20] D. Hoff and K. Zumbrun, *Multi-dimensional diffusion waves for the Navier-Stokes equations of compressible flow*. Indiana Univ. Math. Jour., 44(1995), 603-676.
- [21] T. Kato, *Strong L^p -solutions of the Navier-Stokes equation in $\mathbb{R}^m$ with applications to weak solutions*. Mathematische Zeitschrift, (187)1984, 471-480.
- [22] P.-L. Lions, *Mathematical Topics in Fluid Mechanics*. Vol.2, Compressible models, Oxford University Press, 1998.
- [23] A. Matsumura and T. Nishida, *The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids*. Proc. Japan Acad. Ser. A Math. Sci. 55(1979), 337-342.
- [24] J. Nash, *Le problème de Cauchy pour les équations différentielles d'un fluide général*, Bulletin de la Soc. Math. de France, 90(1962), 487-497.
- [25] M. Paicu and Z. Zhang, *Global Regularity for the Navier-Stokes equations with large, slowly varying initial data in the vertical direction*, preprint, 2009.
- [26] F. Weissler, *The Navier-Stokes Initial Value Problem in L^p* . Arch. Rational Mech. Anal., (74)1980, 219-230.

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS, P.O. Box 8009, BEIJING 100088, P. R. CHINA

E-mail address: chen_qionglei@iapcm.ac.cn

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS, P.O. Box 8009, BEIJING 100088, P. R. CHINA

E-mail address: miao_changxing@iapcm.ac.cn

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, 100871, P. R. CHINA

E-mail address: zfzhang@math.pku.edu.cn